

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/92031/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Unsworth, Richard, Jones, Benjamin and Cullen-Unsworth, Leanne 2016. Seagrass meadows are threatened by expected loss of peatlands in Indonesia. *Global Change Biology* 22 (9) , pp. 2957-2958.
10.1111/gcb.13392

Publishers page: <http://dx.doi.org/10.1111/gcb.13392>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Received Date : 10-May-2016

Revised Date : 31-May-2016

Accepted Date : 01-Jun-2016

Article type : Letters to Editor

Seagrass meadows are threatened by expected loss of peatlands in Indonesia

Richard K.F. Unsworth¹, Benjamin L Jones², Leanne C. Cullen-Unsworth²

¹Seagrass Ecosystem Research Group, College of Science, Swansea University SA2 8PP, UK

²Sustainable Places Research Institute, Cardiff University, 33 Park Place, Cardiff, CF10 3BA, UK

Corresponding Author Email ID: r.k.f.unsworth@swansea.ac.uk

Seagrass meadows provide one of the most productive stores of carbon in our oceans. They also support marine biodiversity and global food security through their role as fish nurseries and fish foraging grounds. Globally their rate of loss is at least as high as that experienced by tropical rainforests. In SE Asia, due to a paucity of long-term data it is difficult to assign such rates of change but significant loss has occurred, possibly up to 40% (Nadiarti *et al.*, 2012, Tomascik *et al.*, 1997). Risks to these meadows continue, with urban development (including coastal development and run-off) being one of the major risks in the region (Grech *et al.*, 2012, Unsworth & Cullen, 2010). Seagrass meadows in Indonesia have also lost their trophic balance due to overexploitation, placing their resilience to poor water quality at risk (Unsworth *et al.*, 2015, Unsworth *et al.*, 2014).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.13392

This article is protected by copyright. All rights reserved.

The article by Abrams et al 2016 utilises a biogeochemical box model to evaluate the downstream effects of the release of Indonesian peat carbon on coastal ecosystems, one of which is seagrass. Their model estimates that the accumulation of detritus in the benthic layer of the coastal environment will lead to an increase in pore water DIN and, therefore to a 31.8% increase in seagrass biomass over the next 60 years (Abrams *et al.*, 2016). This is based on the broad assumption that seagrass in Indonesia is all nutrient limited. In order to reach this conclusion the authors utilise information from just two research papers that examine the mass-balance of nutrients in two Indonesian seagrass meadows conducted in the early 1990's (Erftemeijer & Middelburg, 1995, Erftemeijer *et al.*, 1993).

We applaud the authors for trying to understand the impacts of such a huge problem but feel their conclusions do not reflect the current status and threats to seagrass in Indonesia. We believe that the assumptions in their model and their conclusions could be improved by considering the following:

- 1) Due to rapid population expansion since the early 1990's and increasing loss of rainforest cover coastal water quality in Indonesia is already in decline, peatland loss will add to this. Some seagrass meadows in the country are likely nutrient limited (van Katwijk *et al.*, 2011) but the evidence suggests that these are the minority with threats to seagrass growing (Nadiarti *et al.*, 2012). The seagrass nutrient data used by Abrams et al to develop their model are insufficient to be used as an indication of the current nutrient status of seagrass throughout Indonesia (Erftemeijer & Middelburg, 1995, Erftemeijer *et al.*, 1993). Data on seagrass nutrient condition in Indonesia is limited, but our understanding of nutrient impacts upon seagrasses

Accepted Article

stems from a much bigger broader global literature (Burkholder *et al.*, 2007). We also know that throughout the wider Indo-Pacific region seagrass nutrient levels (from similar species) are commonly elevated (Ambo-Rappe, 2014, McKenzie *et al.*, 2012). An improved model needs to consider this wider body of literature and data.

- 2) Seagrass diversity in Indonesia is high, as a result these seagrasses live across varied environments (estuarine, reef and coastal environments of either an intertidal or subtidal depth) where species have different adaptations. Some reef seagrass meadows in Indonesia far from land will be nutrient poor and therefore increased DIN from peatland degradation will be of benefit resulting in increased growth, but most seagrass meadows (especially coastal and estuarine meadows) in Indonesia are probably already in states of elevated DIN and are at risk. We also know that many short-term experimental enrichment studies on nutrient limited seagrass result in short term increases in biomass but long-term studies find elevated nutrients to result in declining seagrass biomass and density (Cabaço *et al.*, 2013). The model created by Abrams *et al* could be enhanced by classifying seagrass meadows according to habitat type and modelling impacts accordingly utilising recent seagrass nutrient data from throughout the region across varied environments and species (e.g. from the Great Barrier Reef seagrass monitoring program).
- 3) The addition of a detrital layer to seagrass benthos will not just result in increased DIN to the sediments and potentially have a negative impact on light attenuation. The addition of detritus will have a whole stimulatory impact upon the microbial, fungal and detrital feeding community. To expect that the detritus would simply end up as DIN in sediment pore water is at best naïve. Furthermore, a likely additional effect of increased organic detritus is to increase the levels of anoxic sulphide stress

within the sediments with follow on effects upon the seagrass growth and productivity (Marbà *et al.*, 2006). In addition the deposition of organic detritus will also likely bring with it increased sediments, known to alter and seagrass community composition (Terrados *et al.*, 1998).

A large biogeochemical model for how terrestrial degradation impacts the coastal environment in the centre of the Worlds biodiversity has potentially far reaching policy implications. We believe that with further refinement the model by Abrams *et al* could achieve this and determine a more likely response of seagrass to land degradation. The creation of such a model needs to be based on sound scientific knowledge of the response of biota to environmental change. Such a model would therefore be of wide use for the management of the terrestrial and coastal environment in Indonesia and beyond.

References

- Abrams JF, Hohn S, Rixen T, Baum A, Merico A (2016) The impact of Indonesian peatland degradation on downstream marine ecosystems and the global carbon cycle. *Global Change Biology*, **22**, 325-337.
- Ambo-Rappe R (2014) Developing a methodology of bioindication of human-induced effects using seagrass morphological variation in Spermonde Archipelago, South Sulawesi, Indonesia. *Marine Pollution Bulletin*, **86**, 298-303.
- Burkholder JM, Tomasko DA, Touchette BW (2007) Seagrasses and eutrophication. *Journal of Experimental Marine Biology and Ecology*, **350**, 46-72.
- Cabaço S, Apostolaki ET, García-Marín P *et al.* (2013) Effects of nutrient enrichment on seagrass population dynamics: evidence and synthesis from the biomass–density relationships. *Journal of Ecology*, **101**, 1552-1562.
- Erftemeijer PLA, Middelburg JJ (1995) Mass balance constraints on nutrient cycling in tropical seagrass beds. *Aquatic Botany*, **50**, 21-36.
- Erftemeijer PLA, Osinga R, Mars AE (1993) Primary production of seagrass beds in South Sulawesi (Indonesia) - A comparison of habitats, methods and species. *Aquatic Botany*, **46**, 67-90.
- Grech A, Chartrand-Miller K, Erftemeijer P *et al.* (2012) A comparison of threats, vulnerabilities and management approaches in global seagrass bioregions. *Environmental Research Letters*, **7**, 024006.
- Marbà N, Santiago R, Díaz-Almela E, Álvarez E, Duarte CM (2006) Seagrass (*Posidonia oceanica*) vertical growth as an early indicator of fish farm-derived stress. *Estuarine, Coastal and Shelf Science*, **67**, 475-483.

- Mckenzie L, Collier CJ, Waycott M, Unsworth RKF, Yoshida RL, Smith N (2012) Monitoring inshore seagrasses of the GBR and responses to water quality. *Proceedings of the 12th International Coral Reef Symposium*, **15B**: 4.
- Nadiarti A, Riani E, Djuwita I, Budiharsono S, Purbayanto A, Asmus H (2012) Challenging for seagrass management in Indonesia. *Journal of Coastal Development*, **15**, 234-242.
- Terrados J, Duarte CM, Fortes MD *et al.* (1998) Changes in community structure and biomass of seagrass communities along gradients of siltation in SE Asia. *Estuarine, Coastal and Shelf Science*, **46**, 757-768.
- Tomascik T, Mah JA, Nontji A, Moosa KM (1997) *The Ecology of the Indonesian Seas (Part II)*, University of Oxford Press, Periplus Editions (HK) Ltd,.
- Unsworth RKF, Collier CJ, Waycott M, Mckenzie LJ, Cullen-Unsworth LC (2015) A framework for the resilience of seagrass ecosystems. *Maine Pollution Bulletin*, **100**, 34-46.
- Unsworth RKF, Cullen LC (2010) Recognising the necessity for Indo-Pacific seagrass conservation. *Conservation Letters*, **3**, 63-73.
- Unsworth RKF, Hinder SL, Bodger OG, Cullen-Unsworth LC (2014) Food supply depends on seagrass meadows in the coral triangle. *Environmental Research Letters*, **9**, 094005.
- Van Katwijk MM, Van Der Welle MEW, Lucassen ECHET *et al.* (2011) Early warning indicators for river nutrient and sediment loads in tropical seagrass beds: A benchmark from a near-pristine archipelago in Indonesia. *Marine Pollution Bulletin*, **62**, 1512-1520.