

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/92310/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Hickey, Jason E., Prymachuk, Steven and Waterman, Heather 2016. Mental illness research in the Gulf Cooperation Council: a scoping review. *Health Research Policy and Systems* 14 (59) , pp. 1-14.
10.1186/s12961-016-0123-2

Publishers page: <http://dx.doi.org/10.1186/s12961-016-0123-2>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **TITLE PAGE**

2 Title: Mental illness research in the Gulf Cooperation Council: a scoping review

3 Authors:

4 Jason E. Hickey^{1,2}, jehickey@ucalgary.ca

5 Steven Pryjmachuk², steven.pryjmachuk@manchester.ac.uk

6 Heather Waterman³, WatermanH1@cardiff.ac.uk

7 ¹University of Calgary Qatar, PO Box 23133, Doha, Qatar

8 ²University of Manchester, Oxford Rd, Manchester M13 9PL, United Kingdom

9 ³Cardiff University, Cardiff, CF10 3XQ

10 *Correspondence concerning this article should be sent to Jason Hickey, PO Box 23133, Doha, Qatar,
11 +974 44065205, jehickey@ucalgary.ca

12

13

14

1	Table of Contents	
2	Introduction	4
3	Theoretical Framework	6
4	Inclusion Criteria	6
5	Identifying Relevant Studies	7
6	Study Screening	7
7	Data Extraction	8
8	Data Analysis	8
9	Results	9
10	Service Preferences	10
11	Illness: symptomology	12
12	Illness: perceived cause	13
13	Illness: perceived impact	14
14	Recovery: traditional healing	15
15	Recovery: family support	16
16	Recovery: religion	16
17	Discussion	17
18	Strengths and Limitations	20
19	Conclusion	21
20	Competing Interests	22
21	Authors' Contributions	22
22	Authors' Information	22
23		
24	<i>Keywords: Arabs, Mental Disorders; Mental Health; Mental Health Services; Middle East;</i>	
25	<i>Qatar; Scoping Review</i>	

Introduction

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

The Gulf Cooperation Council (GCC) is a union of six Arabic states in the Persian Gulf, including Bahrain, Kuwait, Oman, Qatar, Saudi Arabia and the United Arab Emirates (UAE). These countries share many cultural, political, religious, economic, and geographical similarities [1]. The aim of the union is to promote regional development through coordination, cooperation and integration among member countries [2].

The GCC has benefited substantially in recent decades from large reserves of oil and natural gas. Qatar, for example, is a small, peninsular country that was once mainly utilized by coastal fishermen, pearl divers, and nomadic Bedouin tribes [3]. Now its capital city, Doha, is being transformed into an ultra-modern metropolis and the country has the highest GDP per capita in the world [4]. The UAE provides another example of rapid economic development; in less than ten years it went from being one of the least developed countries in the world to a modern industrialized nation [5].

Islam is the foundation of cultural and social customs in the GCC [6]. It infuses nearly all aspects of life, from architecture, food choice, daily routine, social interactions, education, health care and more. Tolerance, hospitality, and modesty are highly valued [7]. However, social customs also result in strict guidelines for what is considered appropriate behavior in certain circumstances. For example, in Saudi Arabia, it is illegal to publically practice any religion other than Islam, and women are required to wear a gown (i.e., Abaya) and headscarf in public [8]. In other Gulf countries, these rules are often relaxed for foreigners. However, social and cultural norms sometimes clash with ongoing modernization and the influx of foreign workers. Because of this, some feel that traditional values are being threatened [9].

The recent, rapid, changes in the region and the pressures or strains that have been mentioned above have seen mental health and mental illness emerge as priority health concerns for

1 all countries in the GCC. In Kuwait, mental health was identified as one of six strategic priorities
2 through a consultation with the World Health Organization [10]. As part of this agenda, mental
3 health services are being integrated into primary health care, and community and home-based
4 services are being developed. Oman also acknowledges the need to scale up mental health services,
5 particularly by increasing the number of available beds, providing training for primary care workers,
6 and implementing a school health program [11]. In Qatar, a National Mental Health Strategy was
7 recently developed [12]. This strategy focuses on system wide change to reduce stigma, improve
8 treatment seeking, increase availability of resources, scale up the workforce, provide services in a
9 variety of locations, and develop standards and guidelines. Bahrain, Saudi Arabia and the UAE have
10 also emphasized mental health as a national priority and service development is underway in these
11 counties as well [13–15].

12 Research on mental illness from GCC countries suggests that sociocultural factors influence
13 people’s experience of mental illness in the region. For example, causal attribution of mental illness
14 to demons (Jinn) prompt people to seek traditional or religious healers frequently [16]; shame can
15 cause families to impose social isolation on a sick family member [17]; extended family structures
16 can promote increased levels of family support and housing [16]; and, religious (Islamic) influences
17 have been linked to non-Western presentations of illness [18]. Hence, for service planning to be
18 most effective, it cannot necessarily rely on international best practices and evidence from other
19 countries. However, it is widely acknowledged that there is limited local research available to guide
20 contextually-appropriate development of mental health services in the region [11,12].

21 The current article aims to systematically review and synthesize regional literature that
22 reports service user perspectives on mental illness in GCC countries, identify major gaps in the
23 literature, and suggest directions for future research. This information will facilitate the
24 development of mental health services in the GCC. It will also provide information for mental health
25 practitioners in non-GCC countries who provide services for Arabic people.

Theoretical Framework

Arksey and O'Malley's [19] framework was used to develop the review protocol. According to one of the most commonly cited definitions, a scoping study "aims to map *rapidly* the key concepts underpinning a research areas and the main sources and types of evidence available..." [20 as cited by Arksey & O'Malley, 2005, emphasis in original]. Scoping studies tend to be inclusive of a range of research designs regardless of where the research sits on the 'evidence hierarchy' [21,22] and seek to provide greater conceptual clarity [23].

Arksey and O'Malley [19] identify four possible reasons to conduct a scoping study: to examine the extent, range and nature of research activity; to determine the value of undertaking a full systematic review; to summarize and disseminate research finding; and, to identify research gaps in the existing literature. Most reports on scoping studies tend to incorporate a combination of these objectives, and outcomes typically include identification of themes in the literature, gaps that have yet to be addressed, and tangible recommendations for practice and research [24–27]. These characteristics make a scoping study well-suited for the aims of the current review.

Methods

Inclusion Criteria

Articles covering one or more of the following common or clinically relevant illnesses [28,29] were included: mood disorders, alcohol and substance use disorders, schizophrenia, Alzheimer's and other dementias, anxiety disorders, obsessive-compulsive disorder, personality disorders, and phobias. Additionally, studies had to be empirical (i.e., based on observed and measured phenomena and derives knowledge from actual experience rather than from theory or belief), published in English, conducted in the Gulf Cooperation Council (Qatar, Saudi Arabia, Kuwait, Bahrain, United Arab Emirates, Oman). **Only studies that reported subjective data from participants were included because the perspectives and lived experiences of service users are critical for**

1 informing a recovery-oriented understanding of mental illness. Subjective data was defined as
2 opinions/experiences collected directly from participants. Articles not meeting these criteria were
3 excluded from the review.

4 **Identifying Relevant Studies**

5

6 First, several databases (CINAHL, Anthropology Plus, MEDLINE, SocINDEX, PsychInfo,
7 Embase, and NCBI PubMed) were searched. The following search string was developed with the
8 assistance of a librarian: ((MM "Mental Health") OR (MM "Mental Disorders+") OR (MM "Mentally Ill
9 Persons")) AND (Cooperation Council for Arab states of the Gulf OR CCASG OR Gulf Cooperation
10 Council OR GCC OR Qatar* OR Emirat* OR Abu Dhabi* OR Bahrain* OR Kuwait* OR Saudi Arabia* OR
11 Oman*). Second, several regional journals (Arab Journal of Psychiatry, Eastern Mediterranean
12 Health Journal, Avicenna, Journal of Local and Global Health Science, Journal of Local and Global
13 Health Perspectives, QScience Connect, and Qatar Medical Journal) were hand searched. Databases
14 and regional journals were searched from inception to December, 2013. Finally, reference lists of
15 articles identified in the previous two strategies were searched.

16 **Study Screening**

17

18 Figure 1 illustrates the study identification and screening process. The first author screened
19 titles from the initial database results (n=2449) and removed duplicates and irrelevant articles.
20 Inclusion criteria were then applied to full abstracts for the remaining articles (n=655) by the first
21 author. The other two authors (SP, HW) each screened a 5% random sample of abstracts to ensure
22 consistent application of inclusion criteria. Full papers for the articles that passed abstract screening
23 (n=80) were obtained and read in full by two authors. An additional 36 articles were excluded at this
24 stage. Eleven articles were included from the hand search of regional journals and reference lists.

1 Fifty-five articles were included in final data analysis. Table 1 demonstrates how each of these 55
2 studies met inclusion criteria.

3 **Data Extraction**

4

5 The first author (JH) extracted data from all 55 articles using a structured extraction
6 template. This template was developed and piloted by the review team. Two authors (SP, HW) each
7 extracted data from half of the final set, meaning data was independently extracted from each
8 article twice. The two extractions were compared and discrepancies resolved through group
9 discussions between the three authors.

10 [Data were extracted from each study under the following categories:](#) general information
11 (e.g., year, profession of primary author), methodology (e.g., design, study location), sample
12 characteristics (e.g., gender, diagnosis), results (e.g., main outcome, subjective outcome), and
13 discussion (e.g., limitations, conclusions). Strengths and weaknesses of individual studies were
14 assessed and recorded during the extraction process.

15 **Data Analysis**

16

17 Descriptive statistics were compiled to illustrate the “extent, nature and distribution” [19] of
18 literature identified. A thematic analysis of the subjective data was also conducted. This purpose of
19 this analysis was to identify and elaborate on the main concepts addressed by the literature. One of
20 the main criticisms of scoping studies is that there is a lack of transparency and rigour in synthesizing
21 and presenting thematic results. Framework analysis [30] was chosen as an analytic approach in
22 order to address these issues.

23 Framework analysis proceeds through a series of logical steps to reach a narrative summary
24 of the results. Data extraction forms were read and re-read to increase familiarity, then codes

1 were applied to identify key concepts within the data. Once all extraction forms had been coded, the
2 initial codes were reviewed and revised to create a conceptual framework. The conceptual
3 framework was then applied back to the extraction form to assess its fit. When the extraction forms
4 had been recoded according to the conceptual framework, the data was entered into a matrix where
5 results could be examined across themes (columns) or article (rows). The majority of analysis was
6 conducted by one author (JH). However, initial codes and a draft of the conceptual framework was
7 reviewed and discussed by the entire team, and the matrix was reviewed by the team to ensure logic
8 and consistency.

9 **Results**

10 Figure 2 displays the publication timeline. The oldest included study was published in 1975
11 and there are three notable peaks in publication frequency in 1988, 2001-2002, and 2010-2013.
12 There is an upward trend in publications over the entire period.

13 The greatest number of studies were conducted in Saudi Arabia (n=21, 38%) followed by
14 Kuwait (n=18, 33%). Five studies (9%) were conducted in each of Qatar, Bahrain and the United Arab
15 Emirates. One study (2%) was conducted in Oman.

16 Psychiatrists acted as first author on the majority of publications (n=39, 70%) followed by
17 researchers from medicine (specialty unspecified) and nursing, who authored four papers each (7%).
18 Profession of other first authors included, epidemiology (n=2, 4%), psychology (n=2, 4%), the
19 behavioural sciences (n=1, 2%), and pharmacy (n=1, 2%). The profession of the primary author was
20 unspecified in two cases.

21 Figure 3 displays which diagnoses were investigated. The most common target was a mixed
22 sample (n=16, 28%), comprising individuals with a range of diagnoses. This was followed by studies
23 examining exclusively schizophrenia (n=14, 25%) and alcohol or substance use disorders (n=13, 23%).
24 None of the included articles targeted Alzheimer's or other dementias.

1 Table 2 displays sample size by gender and diagnosis. Overall, males outnumber females by a
2 ratio of 2.2 to 1. The largest gender imbalance occurred for alcohol or substance use disorders
3 where males outnumbered females by a ratio of 72.6 to 1. The only disorder where females were
4 over represented was obsessive compulsive disorder, where females outnumbered males by a ratio
5 of 2.6 to 1.

6 The vast majority of studies (n=42, 84%) recruited participants from public psychiatric
7 treatment centers. Of these, half recruited from inpatient or detox units, while half recruited from
8 outpatient departments. Two studies (4%) recruited from private clinics, and only 1 study (2%)
9 recruited participants from the community. Twelve studies (24%) were unclear about recruitment
10 location.

11 Several themes and subthemes emerged during analysis. These include: service preferences,
12 illness (perceived cause, symptomology, impact), and recovery (traditional healing, family support,
13 religion). The following section, which critically synthesizes the results of this review, is organized
14 according to these themes and sub-themes.

15 **Service Preferences.** Gender seemed to influence service preferences in several studies. For
16 example, [Bener and Ghuloum \[31\]](#) found that patients' gender affected the type of topics seen as
17 most important when receiving treatment from a psychiatrist. In that study, males viewed
18 discussions with a doctor about treatment options as being most important, while females
19 prioritized explanations of the condition and the underlying cause. Surveyed preferences were
20 limited to interactions with a psychiatrist; thus, it is not possible to determine preferences across
21 other professional groups. Additionally, the results were collected using a brief questionnaire, which
22 limited choices and did not explore underlying reasons for observed differences. However, the study
23 suggests that gender is an important consideration for service delivery in this region.

1 The importance of gender is reinforced by [Amin and Hamdi \[32\]](#), who found gender to have
2 an influence on where participants preferred to seek treatment. The authors found that females in
3 need of psychiatric care tend to present at the emergency department, while males were more
4 often seen in the outpatient department. [Amin and Hamdi \[32\]](#) suggest several reasons for their
5 observed difference but these were conjecture and not grounded in the results. Additionally, this
6 study is now 20 years old and with recent modernization in the region, results may no longer be
7 accurate. Thus, there has been no recent research to examine service users' preferences across a
8 range of services within the healthcare system.

9 Other studies, however, have demonstrated a preference for services outside of the health
10 care system. [Salem et al. \[33\]](#) found that nearly half of the sample went to a faith healer prior to
11 seeking psychiatric care. The majority of these participants continued to see a faith healer even after
12 engaging psychiatric services. Because convenience sampling was used in this study it is difficult to
13 generalize results to the wider population. However, the participants originated from various
14 countries in the Gulf region, had a range of diagnoses, and included nearly equal proportions of men
15 and women. This diverse sample adds to the generalizability of the study. The author concludes that
16 mental health professionals need to be aware of patient preferences for traditional healing and
17 understand the reasons why they sometimes refuse medical treatment.

18 A comparable study conducted by [Al-Solaim and Loewenthal \[34\]](#) in Saudi Arabia
19 demonstrated that psychiatric services seem to be seen as a last resort when other options (e.g.,
20 faith healers) are not successful. This reinforces [Salem et al.'s \[33\]](#) argument that mental health
21 professionals should not ignore the contribution of traditional healers to service users' treatment.
22 However, [Al-Solaim and Loewenthal's \[34\]](#) results are drawn solely from the experiences of 15
23 women. It is possible that they may not represent the majority view. Men, in particular, as
24 demonstrated earlier, may have different preferences.

1 **Illness: symptomology.** Symptomology was one of the most commonly occurring themes in
2 the included articles. Clinical presentation of mental illness was described for alcohol and substance
3 abuse [35–42], depression [32,43–45], suicide [46,47], panic disorder [48], obsessive-compulsive
4 disorder [34,49,50], schizophrenia [51–55], smoking behavior [56], and hypochondriasis [18]. [Several](#)
5 [of these articles emphasized the unique presentation of certain illness within the Arab context.](#)

6 For example, [Shooka et al. \[50\]](#) demonstrated that religious and blasphemous thoughts were
7 the most common obsessions in patients with obsessive-compulsive disorder (OCD), while repeated
8 prayer-related cleaning and washing was the most common compulsion. [The authors suggest that](#)
9 [there are higher levels of religious content in participants from a strict religious context.](#) These
10 findings were reinforced by [Mahgoub and Abdel-Hafeiz \[49\]](#) who found an [Islamic focus](#) for
11 obsessions and compulsions in OCD. In particular, prayer and body washing were the most common
12 obsessions and religious repeating and religious washing were the most common compulsions. Both
13 these studies are quite old; however, [the Islamic influence](#) on symptomology was also apparent in
14 [Al-Solaim and Loewenthal's \[34\]](#) more recent study on OCD. An additional finding in this more recent
15 was that symptoms in the religious domain were the most disturbing for patients and their families.
16 Despite their limitations, these studies suggest that the content and focus of OCD symptoms are
17 culturally influenced.

18 Research on other disorders has also demonstrated a contextual influence. For example, a
19 study by [Kent and Wahass \[52\]](#) on schizophrenia in Saudi Arabia found that hallucinations had more
20 religious and superstitious content compared to a sample from the UK. [Zarrouk \[55\]](#) found that
21 delusions also differed from non-Arabic samples, with Saudi patients more frequently believing they
22 were being 'made' to do things. In a study on depression by [Hamdi et al. \[44\]](#), four main types of
23 symptom variation (compared to non-Arabic studies) were identified: variations in idioms (e.g.,
24 heavy/tense vs. depressed/sad), use of somatic metaphors (e.g., 'my body is shattered'), influence of

1 religion (e.g., denying acts considered *haram*/forbidden), and behavioral alterations (e.g., going
2 into desert to stare at nothing). Several of these variations clearly reflect the local context.

3 Taken together, these articles demonstrate a contextual influence on symptomology in the
4 GCC. It should be noted that religion is also associated with symptoms in non-Arabic contexts.
5 However, the religious content of symptoms in this review are clearly shaped by the Islamic context.
6 This finding has appeared repeatedly over a considerable period of time (1978-2011), which suggests
7 that this is a somewhat stable phenomenon. Many of these authors reasonably suggest that an
8 understanding of local variations in clinical presentation is important for accurate diagnosis and
9 treatment of service users.

10 **Illness: perceived cause.** The perceived cause of mental illness was often external. For
11 example, participants with a diagnosis of nosophobia attributed the cause to over-investigation of
12 minor complaints by physicians [48]. Similarly, over one third of women in polygamous marriages
13 attributed their illness to their marriage [57], the majority of caregivers attributed the cause of
14 illness to social stressors [58], and those with thoughts of self-harm attributed these thoughts to the
15 devil [48]. Taken individually, the design weaknesses (e.g., convenience sample, bias, cross sectional
16 design) in these articles prevent reliable conclusions from being made. However, the consistent
17 external causal attribution across studies lends credibility to the finding.

18 One particular class of perceived external cause, the supernatural (e.g., black magic, jinn, evil
19 eye), seems particularly common. For example, Salem et al. [33] found that about one third of
20 participants attributed their illness to supernatural factors, while one third attributed them to
21 psychiatric problems and one third were unsure. Similarly, Al-Solaim and Loewenthal [34] and Al-
22 Sughayir [59] found that the majority of patients attributed their illness to possession by jinn (i.e.,
23 evil spirits). This external attribution to supernatural forces can be protective as certain symptoms
24 may not be as stigmatized and may not produce such strong feelings of guilt [34,48]. In fact, those

1 with the evil eye may see themselves as having some positive attribute worthy of envy, which boosts
2 their self-esteem [34]. Again, there are design weaknesses (e.g., bias, unrepresentative samples), but
3 the consistent findings on supernatural attribution warrant further, more rigorous investigation.

4 **Illness: perceived impact.** Most of the data under this theme come from studies on alcohol
5 and substance abuse, which limits applicability to people with other diagnoses. Additionally, there
6 was a certain amount of heterogeneity in impact outcomes. For example, [Al-Ansari and Negrete \[36\]](#)
7 surveyed people undergoing treatment for alcohol abuse. Participants felt guilty about drinking, that
8 it caused their family/friends to worry about them, that it sometimes created interpersonal
9 problems, and caused them to neglect personal responsibilities. In contrast, [Zaidan et al. \[42\]](#) found
10 that the majority of their sample never felt guilty about their drinking. Based on the information
11 presented, the samples in the two studies seem comparable, aside from country of residence.
12 Hence, the reasons for the observed differences are unexplained. Additionally, both studies enrolled
13 males only so results are further limited.

14 [Daradkeh and Karim \[60\]](#) and [Al-Solaim and Loewenthal \[34\]](#) offer limited information on
15 illness impact among people with schizophrenia and OCD, respectively. These included barriers to
16 social inclusion (schizophrenia) and interpersonal problems with family members (schizophrenia and
17 OCD). However, the small sample size used in these studies means that further investigation is
18 needed before reliable conclusions can be made.

19 Another major limitation in the majority of the studies on alcohol and substance abuse is
20 their almost exclusive reliance on questionnaires to collect data. This limits the depth and richness of
21 information collected and prevents participants from sharing relevant information that is not
22 covered on the questionnaire. Thus, despite the relatively large number of studies on this topic,
23 understanding remains limited.

1 **Recovery: traditional healing.** As mentioned above, a large proportion of psychiatric
2 patients visit a traditional healer prior to seeking medical help [32–34,58,61]. This practice was
3 perceived to contribute to recovery in a variety of ways. However, there were mixed perceptions on
4 the effectiveness of traditional treatment.

5 Qureshi et al. [62] found that some participants with depressive or catatonic symptoms
6 reported a temporary improvement from traditional treatment but that most were unsatisfied.
7 Similarly, Salem et al. [33] found that about half of participants with a range of diagnoses
8 experienced only a temporary benefit, with others experiences no benefit at all. Both studies
9 interpreted ‘benefit’ as a reduction in symptoms.

10 While most authors exploring the effectiveness of traditional healing seem to assume that
11 symptom reduction is of primary importance, this may not necessarily be the case for service users.
12 For example, Al-Subaie [61] found that even those who did not perceive their symptoms to be
13 reduced reported feeling that God would reward them for having faith in traditional healing
14 methods, which are primarily based on religious beliefs. This finding suggests that service users may
15 place value on treatment benefits other than a reduction of symptoms. However, these other
16 potential benefits are largely ignored in the studies that were reviewed, as is the relative importance
17 of various benefits to participants.

18 A further limitation in the studies included in this section is selection bias. Since participants
19 were selected from the population of people seeking psychiatric treatment, it is possible that others
20 received longer lasting benefit from traditional treatments and did not subsequently access services.
21 A broader sampling strategy (e.g., including people with mental illness who do not make regular use
22 of psychiatric services) would be necessary for a more accurate investigation.

1 **Recovery: family support.** The extended nature of the Gulf Arab family was frequently
2 addressed as being a source of support for service users. Potential supportive roles of the family in
3 one study included medication supervision, being tolerant of short periods of withdrawal, helping to
4 find acceptable ways to describe and understand the illness, not expecting anything in return for
5 their help, and assistance in filling leisure time [63]. While these caregiving themes make sense
6 intuitively, the author introduces subjectivity and bias into the analysis and does not consider
7 alternate or contradictory views. Additionally, the study was conducted over 20 years ago and no
8 similar studies have been conducted to support or refute the author's findings. It is also possible that
9 families' supportive roles have changed in the ensuing period of rapid socioeconomic development.

10 Studies also suggested that support is more common in extended families compared to
11 nuclear families and that those living in extended families had more social contact, better personal
12 hygiene, less active symptoms, and better treatment outcomes [63,64]. Conversely, the extended
13 family was also perceived to have a negative impact; tension or stress within the family were cited as
14 reasons for substance abuse [65], attempted suicide [47] and were associated with higher disease
15 severity [53]. Unfortunately, only [Zahid and Ohaeri's \[53\]](#) study was recent enough to consider these
16 influences within a modern context, and even this study did not examine the issue in depth. Hence,
17 while it seems likely that the family plays a role in service users' recovery from mental illness, the
18 nature of this role in a modern context remains unclear.

19 **Recovery: religion.** Committing acts that were incongruent with the teachings of Islam led to
20 feelings of guilt and lower self-esteem [34,48]. The more compliant a person was with the values of
21 their faith, the more pride they felt [34]. However, compulsive religious acts relating to mental
22 illness sometimes interfered with daily life and led to treatment seeking. Religion however,
23 particularly prayer, was still seen as one of the main ways to cope with mental illness and related

1 stress. For example, being religious was associated with lower levels of death anxiety [39] and
2 alcohol abuse [64,66].

3 The majority of studies included in this section incorporated religion as a minor variable that
4 was a small part of a larger study. Thus, a systematic, in-depth investigation of religion has not been
5 undertaken. Despite a lack of direct evidence, many of the reviewed studies argued that religion had
6 an impact on certain aspects of participants' illness, including symptomology [e.g., ,43], treatment
7 seeking [e.g., 37], etc. However, these claims and their underlying assumptions are currently
8 unsubstantiated by the literature. In other words, broad assumptions were sometimes made, based
9 solely on the authors' own personal beliefs or professional understanding about the value of
10 religion. Thus, while it seems likely that religion play a role, it is difficult to objectively interpret the
11 impact of religion based on the literature included in this review.

12 Discussion

13

14 This systematic scoping review was undertaken in order to synthesize regional literature on
15 service users' experience of mental illness, to identify gaps in the literature and identify
16 opportunities for future research. Fifty-five articles were included in the review.

17 The included studies offer a small glimpse into service preferences, including a preference
18 for initial consultations with faith healers. The use of faith healers to treat mental illness has also
19 been documented in other developing countries. For example, a study conducted in rural India
20 reported that faith healing is widely used and that many people seek traditional services before
21 medical psychiatric services [67]. Services of faith healers are also commonly used in Ghana. This is
22 because traditional healers offer more culturally appropriate models of understanding illness, higher
23 levels of psychosocial support, and easier accessibility [68]. A study in Zimbabwe found that three

1 quarters of people sought both traditional and medical treatment for mental illness; however, in this
2 study, biomedical care providers were the most common point of first contact [69].

3 The frequent use of traditional healers has implications for government-run psychiatric
4 services. The title of the article by [Ae-Ngibise et al. \[68\]](#) summarizes these implications quite well:
5 ‘Whether you like it or not, people with mental health problems are going to go to [faith healers]’.
6 While some collaboration currently occurs in Qatar, ongoing service development efforts should aim
7 to identify opportunities to incorporate safe, appropriate traditional healing as part of a
8 comprehensive service.

9 External stressors, particularly the supernatural (i.e., jinn, black magic, evil eye) were
10 frequently seen as the root cause of mental illness. This indicates a need to address cultural beliefs
11 and social factors when treating mental illness. A recently published review found that psychiatric
12 symptoms are commonly attributed to the supernatural among Muslims worldwide [70]. The
13 authors claim that this external attribution has diagnostic and treatment implications. For example,
14 biomedical treatment may not be accepted if underlying cultural beliefs about the supernatural are
15 not addressed [71]. Therefore, practitioners should foster an awareness of traditional beliefs and be
16 open to incorporating these as part of the therapeutic process.

17 Symptomology was described for most major mental illnesses and several similarities and
18 differences were identified compared to typical presentations in the West. Variations in
19 presentation of mental illness across cultures has been widely reported and debated in the literature
20 [72]. Variations create difficulties in applying standard diagnostic criteria such as the DSM-5 or the
21 ICD-10 in cultural contexts that differ from the West. This provides fuel for an argument that the
22 dichotomous nature of diagnostic categories are unhelpful in the treatment of people with mental
23 health issues [73,74]. Unlike physical illness, mental illness is predominantly subjective (i.e.,
24 unmeasurable); interpretation and explanation of symptoms by the patient is influenced by

1 sociocultural understanding [75,76]. Attending to a person's subjective experience of their illness
2 and the overall impact of the illness on their life can lead to a more holistic understanding of the
3 patient and better treatment outcomes [77]. It is worth noting that this practice aligns well with
4 recovery-oriented care.

5 Other sociodemographic and sociocultural factors such as family support, marriage, religion,
6 education, financial status, gender, birth order, and nationality/ethnicity were also investigated.
7 However, the relationship of these factors to mental illness was difficult to assess due to limited data
8 and weaknesses of the included studies. These areas provide fertile ground for future research.

9 The articles included in this review add important, contextually-relevant, data to our
10 understanding of mental illness in the region. However, despite nearly 40 years of research
11 addressing the impact of sociocultural factors on people's experiences of mental illness, a limited
12 understanding of these issues remain. For example, the Arab extended family is widely
13 acknowledged in having a key supportive role for those with mental illness. However, it has also
14 been reported as a source of stress and conflict. Religion influences all aspects of life in Arab society
15 and provides a source of strength and support for people with mental illness. However, forbidden
16 acts such as alcohol and substance abuse and suicide still occur, and cause guilt and worry for
17 patients and their families. Widespread belief in the influence of the supernatural on mental illness
18 has been documented. These beliefs fall within the scope of modern mental health practice, but
19 there is little evidence of how they can be incorporated successfully into a contextually-relevant
20 model of psychiatric care and recovery. Research into the interplay between psychiatry and
21 traditional healing is also lacking; service users place value on consultation with faith healers, but it
22 is unclear how these services might be incorporated into a cohesive system of mental health service
23 in the region, or even if they should be incorporated. The concept of stigma is conspicuously absent
24 from the articles and little effort has gone into measuring or describing stigma towards mental

1 illness in the GCC. Also absent from the literature are investigations into service users' self-
2 management of their illness. Most service users spend the majority of their lives outside of the
3 health care system. Yet, virtually nothing is known about the strategies and resources they use to
4 minimize the day-to-day challenges of mental illness.

5 **Strengths and Limitations**

6

7 This review is the first of its kind to be conducted in the Gulf region. Two other reviews of
8 mental health research have been published [78,79]. However, the scope of these reviews is limited
9 to the frequency, distribution and topics of publications. This review is the first to synthesize the
10 results of research on mental illness in this region. Additionally, the systematic approach undertaken
11 for this review facilitated a relatively objective synthesis that was rigorously conducted. Finally, the
12 use of framework analysis overcomes a major criticism of scoping studies by providing a transparent
13 method of analysis where thematic results can be clearly linked to individual studies.

14 This review is limited by the general weaknesses in the body of literature. For example, most
15 of the research included in this review seems to be unfunded. This implies that proposals may not
16 have been subjected to peer review. *While lack of funding does not necessarily imply low quality,*
17 *many of the papers reviewed here would have benefitted from additional peer review during the*
18 *planning stage.* Increasing the funding available for mental health research could help to improve
19 the overall quality of the research (e.g., through the scientific review process) and guide the focus of
20 future research to ensure it is ethically sound and relevant for the development of practice and
21 policy.

22 The evidence base consists primarily of cross-sectional studies aimed at developing
23 foundational knowledge. This type of evidence does not allow for causal inferences to be made and
24 only provides a snapshot of a phenomenon at one point in time. This means that the relationships

1 between sociodemographic factors and illness experiences are still not well understood. Longitudinal
2 research would help to increase understanding of illness trajectory outside of the hospital setting,
3 while interventional research would help to improve the transformation of knowledge into practice.
4 Additionally, the vast majority of included research was quantitative, meaning that results of the
5 review lack depth. Incorporating more qualitative research has the potential to clarify important
6 issues, help to develop a better understanding of service users' perspectives, and build contextually
7 relevant mental health theory that can be applied and tested through subsequent research.
8 Qualitative research would also facilitate development of contextually valid measurement scales and
9 questionnaires.

10 Underrepresentation of women in the studies highlights the need to look beyond
11 convenience samples in psychiatric research at ways of identifying more representative samples. The
12 gender imbalance implies that review findings are more relevant to the male psychiatric population.
13 Finally, very few studies include family members or caregivers, even though most acknowledge the
14 contribution of social support to recovery. Thus, understanding of caregivers' potentially supportive
15 role in the region remains limited.

16 **Conclusion**

17

18 Despite the limitations of the reviewed literature, we can conclude that the sociocultural
19 context in the Gulf region is linked to people's experience of mental illness. However, service users'
20 perceptions and understandings about the nature of the context-illness experience relationship have
21 not been systematically explored. This is particularly true for the process of self-management of
22 illness outside the hospital setting. Interventions that are developed based on this limited
23 understanding may have limited effectiveness and acceptability. While many questions can be
24 derived from the identified knowledge gaps, two seem prominent:

1 1. What strategies do service users in Arabic countries use to self-manage their mental illness
2 outside of the hospital setting?

3 2. What treatments and interventions would be most effective and acceptable to support this
4 self-management?

5 Future research that attempts to address these two questions will increase the capacity of Arabic
6 mental health services to provide efficient and culturally appropriate support to service users.

7 **Competing Interests**

8
9 The authors declare that they have no competing interests.

10 **Authors' Contributions**

11

12 All authors made substantial contributions to the conception and design of this study. JH was
13 responsible for article retrieval, study screening, data extraction and analysis, and manuscript
14 preparation. SP and HW participated in study screening, data extraction and analysis, and critically
15 review and revision of the manuscript. All authors read and reviewed the final manuscript.

16 **Authors' Information**

17

18 JH has been working as a Nursing Instructor in Qatar since 2012. His research focuses on service
19 users' self-management of mental illness in the Middle East. This article was undertaken as a part of
20 JH's PhD studies at the University of Manchester.

21

22

1 References

- 2 1. AlMarri TSK, Oei TPS. Alcohol and substance use in the Arabian Gulf region: A review. *International*
3 *Journal of Psychology*. 2009;44:222–33.
- 4 2. Cooperation Council for the Arab States of the Gulf. Cooperation Council for the Arab States of the
5 Gulf. The charter [Internet]. Cooperation Council for the Arab States of the Gulf; 2012. Available from:
6 <https://www.gcc-sg.org/eng/indexfc7a.html?action=Sec-Show&ID=1>
- 7 3. Kronfol Z, Ghuloum S, Weber A. Country in focus: Qatar. *Asian J Psychiatr*. 2013;6:275–7.
- 8 4. Knoema. GDP per Capita Ranking 2015 [Internet]. Knoema. 2015 [cited 2016 Jun 9]. Available from:
9 <http://knoema.com/sijweyg/gdp-per-capita-ranking-2015-data-and-charts>
- 10 5. Shihab M. Economic Development in the UAE. *United Arab Emirates: a new perspective*. 2001;249–
11 259.
- 12 6. Williams J. Don't They Know It's Friday? Cross-Cultural Considerations for Business and Life in the Gulf.
13 1ST edition. London: Motivate Publishing; 1999.
- 14 7. Government of Dubai Executive Council. The Dubai Code of Conduct [Internet]. n.d. Available from:
15 http://www.zu.ac.ae/employment/html/documents/CultureandConductinDubai_000.pdf
- 16 8. Saudi Arabia. Local laws and customs [Internet]. 2016 [cited 2016 Apr 10]. Available from:
17 <https://www.gov.uk/foreign-travel-advice/saudi-arabia/local-laws-and-customs>
- 18 9. Fellow J. Qatar's delicate balancing act [Internet]. BBC News. 2013. Available from:
19 <http://www.bbc.co.uk/news/world-middle-east-21029018>
- 20 10. World Health Organization. Country Cooperation Strategy for WHO and Kuwait 2012–2016 [Internet].
21 Cairo: WHO Regional office for the Eastern Mediterranean; 2014. Available from:
22 <http://apps.who.int/iris/handle/10665/113231>
- 23 11. World Health Organization. Country Cooperation Strategy for WHO and Oman 2010-2015 [Internet].
24 Cairo: WHO Regional office for the Eastern Mediterranean; 2010. Available from:
25 http://www.who.int/countryfocus/cooperation_strategy/ccs_omn_en.pdf
- 26 12. Supreme Council of Health. Qatar National Mental Health Strategy. *Changing Minds, Changing Lives:*
27 *2013-2018* [Internet]. 2013 [cited 2013 Dec 15]. Available from: <http://www.nhsq.info/app/media/1166>
- 28 13. Qureshi NA, Al-Habeeb AA, Koenig HG. Mental health system in Saudi Arabia: an overview.
29 *Neuropsychiatr Dis Treat*. 2013;9:1121–35.
- 30 14. Kingdom of Bahrain Ministry of Health. Bahrain Health Strategy. *Framework for Action* [Internet].
31 Bahrain: Ministry of Health; 2002. Available from:
32 <http://www.moh.gov.bh/pdf/bahrainhealthstrategy.pdf>

- 1 15. World Health Organization. Country cooperation strategy for WHO and United Arab Emirates 2012-
2 2017 [Internet]. Cairo: WHO Regional office for the Eastern Mediterranean; 2012. Available from:
3 <http://apps.who.int/iris/handle/10665/113226>
- 4 16. El-Islam M. Arab Culture and Mental Health Care. *Transcultural Psychiatry*. 2008;45:671–82.
- 5 17. Al-Krenawi A, Graham JR. Culturally sensitive social work practice with Arab clients in mental health
6 settings. *Health Soc Work*. 2000;25:9–22.
- 7 18. El-Islam MF, Moussa MA, Malasi TH, Suleiman MA, Mirza IA. Assessment of depression in Kuwait by
8 principal component analysis. *Journal of Affective Disorders*. 1988;14:109–14.
- 9 19. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International Journal of*
10 *Social Research Methodology*. 2005;8:19–32.
- 11 20. Mays N, Pope C, Popay J. Systematically reviewing qualitative and quantitative evidence to inform
12 management and policy-making in the health field. *J Health Serv Res Policy*. 2005;10:6–20.
- 13 21. Akers J. *Systematic reviews: CRD's guidance for undertaking reviews in health care*. York: CRD,
14 University of York; 2009.
- 15 22. Anderson S, Allen P, Peckham S, Goodwin N. Asking the right questions: Scoping studies in the
16 commissioning of research on the organisation and delivery of health services. *Health Res Policy Syst*.
17 2008;6:7.
- 18 23. Davis K, Drey N, Gould D. What are scoping studies? A review of the nursing literature. *International*
19 *Journal of Nursing Studies*. 2009;46:1386–400.
- 20 24. Han CS, Oliffe JL, Ogrodniczuk JS. Suicide among East Asians in North America: A scoping review.
21 *Journal of Mental Health*. 2013;22:361–71.
- 22 25. Jenkins E, Goldner EM. Approaches to Understanding and Addressing Treatment-Resistant
23 Depression: A Scoping Review. *Depress Res Treat* [Internet]. 2012 [cited 2013 Oct 31];2012. Available
24 from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3337614/>
- 25 26. Ross F, Smith E, Mackenzie A, Masterson A. Identifying research priorities in nursing and midwifery
26 service delivery and organisation: a scoping study. *International Journal of Nursing Studies*. 2004;41:547–
27 58.
- 28 27. Saunders SL, Nedelec B. What Work Means to People with Work Disability: A Scoping Review. *J Occup*
29 *Rehabil*. 2013;1–11.
- 30 28. World Health Organization. *The Global Burden of Disease: 2004 Update* [Internet]. Geneva; 2008.
31 Available from:
32 http://www.who.int/healthinfo/global_burden_disease/GBD_report_2004update_full.pdf

- 1 29. Ghuloum S, Bener A, Abou-Saleh MT. Prevalence of mental disorders in adult population attending
2 primary health care setting in Qatari population. *J Pak Med Assoc.* 2011;61:216–21.
- 3 30. Ritchie J, Lewis J. *Qualitative research practice: a guide for social science students and researchers.*
4 London; Thousand Oaks, Calif.: Sage Publications; 2003.
- 5 31. Bener A, Ghuloum S. Gender difference on patients' satisfaction and expectation towards mental
6 health care. *Niger J Clin Pract.* 2013;16:285–91.
- 7 32. Amin Y, Hamdi E. Gender differences in psychiatric admissions: The influence of culture and social
8 structure. *Arab Journal of Psychiatry.* 1995;6:66–81.
- 9 33. Salem MO, Saleh B, Yousef S, Sabri S. Help-seeking behaviour of patients attending the psychiatric
10 service in a sample of United Arab Emirates population. *Int J Soc Psychiatry.* 2009;55:141–8.
- 11 34. Al-Solaim L, Loewenthal K. Religion and obsessive-compulsive disorder (OCD) among young Muslim
12 women in Saudi Arabia. *Mental Health, Religion & Culture.* 2011;14:169–82.
- 13 35. Abalkhail B. Social status, health status and therapy response in heroin addicts. *Eastern*
14 *Mediterranean Health Journal.* 2001;7:465–72.
- 15 36. Al-Ansari EA, Negrete JC. Screening for alcoholism among alcohol users in a traditional Arab Muslim
16 society. *Acta Psychiatrica Scandinavica.* 1990;81:284–8.
- 17 37. Al-Kandari F, Yacoub K, Omu F. Effect of drug addiction on the biopsychosocial aspects of persons
18 with addiction in Kuwait: nursing implications. *Journal of Addictions Nursing.* 2007;18:31–40.
- 19 38. Bilal AM, Kristof J, El-Islam MF. A cross-cultural application of a drinking behaviour questionnaire.
20 *Addictive Behaviors.* 1987;12:95–101.
- 21 39. Daradkeh F, Moselhy HF. Death anxiety (Thanatophobia) among drug dependents in an Arabic
22 psychiatric hospital. *American Journal of Drug & Alcohol Abuse.* 2011;37:184–8.
- 23 40. Iqbal N. Recoverable hearing loss with amphetamines and other drugs. *Journal of Psychoactive Drugs.*
24 2002;36:285–8.
- 25 41. Qureshi NA. Trihexyphenidyl (artane) abuse among Saudi psychiatric patients. *Ann Saudi Med.*
26 1992;12:391–4.
- 27 42. Zaidan ZAJ, Dorvlo ASS, Viernes N, Al-Suleimani A, Al-Adawi S. Hazardous and harmful alcohol
28 consumption among non-psychotic psychiatric clinic attendees in Oman. *International Journal of Mental*
29 *Health and Addiction.* 2006;5:3–15.
- 30 43. El-Islam MF, Moussa MA, Malasi TH, Suleiman MA, Mirza IA. Assessment of depression in Kuwait by
31 principal component analysis. *Journal Of Affective Disorders.* 1988;14:109–14.

- 1 44. Hamdi E, Amin Y, Abou-Saleh MT. Problems in validating endogenous depression in the Arab culture
2 by contemporary diagnostic criteria. *Journal Of Affective Disorders*. 1997;44:131–43.
- 3 45. Suleiman RA, Khashaba AS, Al-Haddad M. Depressive symptoms among HIV positive drug users in
4 Bahrain. *Arab Journal of Psychiatry*. 2002;1.
- 5 46. Al-Habeeb AA, Sherra KS, Al-Sharqi AM, Qureshi NA. Assessment of suicidal and self-injurious
6 behaviours among patients with depression. *East. Mediterr. Health J*. 2013;19:248–54.
- 7 47. Suleiman MA, Nashef AA, Moussa MAA, El-Islam MF. Psychosocial profile of the parasuicidal patient
8 in Kuwait. *International Journal of Social Psychiatry*. 1986;32:16–22.
- 9 48. El-Islam MF. Cultural aspects of morbid fears in Qatari women. *Social Psychiatry and Psychiatric
10 Epidemiology*. 1994;29:137–40.
- 11 49. Mahgoub OM, Abdel-Hafeiz HB. Pattern of obsessive-compulsive disorder in eastern Saudi Arabia.
12 *The British Journal Of Psychiatry: The Journal Of Mental Science*. 1991;158:840–2.
- 13 50. Shooka A, al-Haddad MK, Raees A. OCD in Bahrain: a phenomenological profile. *The International
14 Journal Of Social Psychiatry*. 1998;44:147–54.
- 15 51. Al-Ansari EA, Emara MM, Mirza IA, El-Islam MF. Schizophrenia in ICD-10: a field trial of suggested
16 diagnostic guidelines. *Comprehensive Psychiatry*. 1989;30:416–9.
- 17 52. Kent G, Wahass S. The content and characteristics of auditory hallucinations in Saudi Arabia and the
18 UK: a cross-cultural comparison. *Acta Psychiatrica Scandinavica*. 1996;94:433–7.
- 19 53. Zahid MA, Ohaeri JU. Schizophrenia psychopathology in a Kuwaiti Arab sample. *Psychopathology*.
20 2010;43:345–56.
- 21 54. Zarrouk ET. The frequency of visual hallucinations in schizophrenic patients in Saudi Arabia. *The
22 British Journal of Psychiatry*. 1975;127:553–5.
- 23 55. Zarrouk ET. The usefulness of first-rank symptoms in the diagnosis of schizophrenia in a Saudi Arabian
24 population. *The British Journal of Psychiatry*. 1978;132:571–3.
- 25 56. Al-Habeeb T, Qureshi N. Smoking among male psychiatric outpatients in Saudi Arabia. *Annals of Saudi
26 Medicine*. 2000;20:218–23.
- 27 57. Chaleby K. Women of polygamous marriages in an inpatient psychiatric service in Kuwait. *J. Nerv.
28 Ment. Dis*. 1985;173:56–8.
- 29 58. Al-Faraj A, Al-Ansari A. What family caregivers of schizophrenic and non-psychotic in-patients in
30 Bahrain believe about mental illness. *Arab Journal of Psychiatry*. 2002;13.

- 1 59. Al Sughayir M. Dissociative experience in psychiatric out patients who have possession belief. Arab
2 Journal of Psychiatry. 2005;16:98–106.
- 3 60. Daradkeh TK, Karim L. Predictors of employment status of treated patients with DSM-III-R diagnosis.
4 Can logistic regression model find a solution? The International Journal of Social Psychiatry. 1994;40:141–
5 9.
- 6 61. Al-Subaie A. Traditional healing experiences in patients attending a university outpatient clinic. Arab
7 Journal of Psychiatry. 1994;5.
- 8 62. Qureshi NA, Al-Amri AH, Abdelgadir MH, El-Haraka EA. Traditional cauterly among psychiatric patients
9 in Saudi Arabia. Transcultural Psychiatry. 1998;35:75–83.
- 10 63. El-Islam MF. Rehabilitation of schizophrenics by the extended family. Acta Psychiatrica Scandinavica.
11 1982;65:112–9.
- 12 64. Bilal AM, Kristof J, Shaltout A, El-Islam MF. Treatment of alcoholism in Kuwait: a prospective follow-up
13 study. Drug and Alcohol Dependence. 1987;19:131–44.
- 14 65. Al-Nahedh N. Relapse among substance-abuse patients in Riyadh, Saudi Arabia. Eastern
15 Mediterranean Health Journal. 1999;5:241–6.
- 16 66. Demerdash AM, Mizaal H, El Farouki S, El Mossalem H. Some behavioural and psychological aspects of
17 alcohol and drug dependence in Kuwait Psychiatric Hospital. Acta Psychiatrica Scandinavica.
18 1981;63:173–85.
- 19 67. Schoonover J, Lipkin S, Javid M, Rosen A, Solanki M, Shah S, et al. Perceptions of traditional healing
20 for mental illness in rural Gujarat. Annals Of Global Health. 2014;80:96–102.
- 21 68. Ae-Ngibise K, Cooper S, Adiibokah E, Akpalu B, Lund C, Doku V, et al. “Whether you like it or not
22 people with mental problems are going to go to them”: a qualitative exploration into the widespread use
23 of traditional and faith healers in the provision of mental health care in Ghana. Int Rev Psychiatry.
24 2010;22:558–67.
- 25 69. Patel V, Simunyu E, Gwanzura F. The pathways to primary mental health care in high-density suburbs
26 in Harare, Zimbabwe. Soc Psychiatry Psychiatr Epidemiol. 1997;32:97–103.
- 27 70. Lim A, Hoek HW, Blom JD. The attribution of psychotic symptoms to jinn in Islamic patients.
28 Transcultural Psychiatry. 2015;52:18–32.
- 29 71. Hanely J, Brown A. Cultural variations in interpretation of postnatal illness: jinn possession amongst
30 Muslim communities. Community Ment Health J. 2013;50:348–53.
- 31 72. Kirmayer LJ, others. Cultural variations in the clinical presentation of depression and anxiety:
32 implications for diagnosis and treatment. Journal of Clinical Psychiatry. 2001;62:22–30.

- 1 73. Slade M. *Personal Recovery and Mental Illness: A Guide for Mental Health Professionals*. 1st edition.
2 Cambridge; New York: Cambridge University Press; 2009.
- 3 74. Uher R, Rutter M. Basing psychiatric classification on scientific foundation: Problems and prospects.
4 *Int Rev Psychiatry*. 2012;24:591–605.
- 5 75. Department of Health and Human Services. *Mental health: Culture, race, and ethnicity. Supplement
6 to mental health: a report of the Surgeon General*. Washington (DC): Government Printing Office
7 [Internet]. U.S. Public Health Service; 2001. Available from:
8 http://www.ncbi.nlm.nih.gov/books/NBK44243/pdf/Bookshelf_NBK44243.pdf
- 9 76. Rosendal M, Jarbøl DE, Pedersen AF, Andersen RS. Multiple perspectives on symptom interpretation
10 in primary care research. *BMC Family Practice*. 2013;14:167.
- 11 77. Kleinman A. *Rethinking Psychiatry: From Cultural Category to Personal Experience*. 1st edition. New
12 York; London: Free Press; 1991.
- 13 78. Afifi MM. Mental health publications from the Arab world cited in PubMed, 1987-2002. *East.
14 Mediterr. Health J*. 2005;11:319–28.
- 15 79. Osman OT, Afifi M. Troubled minds in the Gulf: mental health research in the United Arab Emirates
16 (1989-2008). *Asia Pac J Public Health*. 2010;22:48S–53S.
- 17 80. Derbas A, al-Haddad M. Factors associated with immediate relapse among Bahraini heroin abusers.
18 *Eastern Mediterranean Health Journal*. 2001;7:473–80.
- 19 81. Al-Kandari F, Yacoub K, Omu F. Initiation factors for substance abuse. *Journal of Advanced Nursing*.
20 2001;34:78–85.
- 21 82. Al-Saffar N, Abdulkareem A, Abdulhakeem A, Salah A-Q, Heba M. Depressed patients' preferences for
22 education about medications by pharmacists in Kuwait. *Patient Educ Couns*. 2008;72:94–101.
- 23 83. Bilal AM. Correlates of addiction-related problems in Kuwait: a cross-cultural view. *Acta Psychiatrica
24 Scandinavica*. 1988;78:414–6.
- 25 84. El-Islam MF, Malasi TA, Suleiman MA, Mirza IA. The correlates of hypochondriasis in depressed
26 patients. *International Journal Of Psychiatry In Medicine*. 1988;18:253–61.
- 27 85. Suleiman MA, Malasi TH, Mirza IA, el-Islam MF. Some characteristics of the psychiatric population
28 attending a primary care centre in Kuwait. *Acta Psychiatr Scand*. 1989;79:199–204.
- 29 86. Zahid M, Ohaeri J, Elshazly A, Basiouny M, Hamoda H, Varghese R. Correlates of quality of life in an
30 Arab schizophrenia sample. *Social Psychiatry & Psychiatric Epidemiology*. 2010;45:875–87.
- 31 87. Zahid M, Ohaeri JU. Relationship of family caregiver burden with quality of care and psychopathology
32 in a sample of Arab subjects with schizophrenia. *BMC Psychiatry*. 2010;10:71–71.

- 1 88. Zahid M, Ohaeri JU. Clinical and psychosocial factors associated with needs for care: an Arab
2 experience with a sample of treated community-dwelling persons with schizophrenia. *Soc Psychiatry*
3 *Psychiatr Epidemiol.* 2013;48:313–23.
- 4 89. Zahid M, Ohaeri JU, Al-Zayed AA. Factors associated with hospital service satisfaction in a sample of
5 Arab subjects with schizophrenia. *BMC Health Serv Res.* 2010;10:294.
- 6 90. Bener A, Al-Kazaz M, Ftouni D, Al-Harthy M, Dafeeah EE. Diagnostic overlap of depressive, anxiety,
7 stress and somatoform disorders in primary care. *Asia Pac Psychiatry.* 2013;5:E29-38.
- 8 91. Ghuloum S, Bener A, Burgut F. Ethnic differences in satisfaction with mental health services among
9 psychiatry patients. *Open Psychiatry Journal.* 2010;4.
- 10 92. Chaleby K. Psychosocial stresses and psychiatric disorders in an outpatient population in Saudi Arabia.
11 *Acta Psychiatrica Scandinavica.* 1986;73:147–51.
- 12 93. Chaleby KS. Perceived parental attitude in social phobia patients. *Ara.* 1990;1.
- 13 94. Shahin SH, Daly EB. Knowledge, attitudes and beliefs about psychotropic medication among Saudi
14 hospitalized psychiatric patients. *Int J Nurs Stud.* 1999;36:51–5.
- 15 95. Al-Sughayir MA. In-patient treatment for resistant obsessive-compulsive disorder. *Saudi Medical*
16 *Journal.* 2000;21:193–5.
- 17 96. Al-Showkan A, Curtis J, White Y. Self-reported quality of life for people with schizophrenia in a
18 psychiatric outpatient department in Saudi Arabia. *Arab Journal of Psychiatry.* 2013;24:93–101.
- 19 97. Chaleby K. Traditional Arabian marriages and mental health in a group of outpatient Saudis. *Acta*
20 *Psychiatrica Scandinavica.* 1988;77:139–42.
- 21 98. Daradkeh TK, Al Habeeb T. Quality of life of patients with schizophrenia 2. *Eastern Mediterranean*
22 *Health Journal.* 2005;11:898–904.
- 23 99. Al-Zarrad F, Abu-Mugaiseeb AM. Abu-Dhabi psychiatric day hospital efficacy and acceptability of its
24 health service (a prospective study). *Arab Journal of Psychiatry.* 2002;13.

1 **Tables and Figures**

2 Table 1. Inclusion data for all articles included in final review

3 Table 2. Gender of participants for each of the studied diagnostic categories.

4 Figure 1. Strategy used to identify studies

5 Figure 2. Number of publications by year of publication.

6 Figure 3. Number of publications by target diagnosis.

7

8

9