

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/96000/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Malmedy, Florence and Wirth, Thomas 2016. Frontispiece: stereoselective ketone rearrangements with hypervalent iodine reagents. *Chemistry - a European Journal* 22 (45) , pp. 16072-16077.
10.1002/chem.201684562

Publishers page: <http://dx.doi.org/10.1002/chem.201684562>

Please note:


Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Stereoselective Ketone Rearrangements with Hypervalent Iodine Reagents

Florence Malmedy and Thomas Wirth*




Abstract: The first stereoselective version of an iodine(III)-mediated rearrangement of arylketones in the presence of orthoesters is described. The reaction products, α -arylated esters, are very useful intermediates in the synthesis of bio-

active compounds such as ibuprofen. With chiral lactic acid-based iodine(III) reagents product selectivities of up to 73% ee have been achieved.

Introduction

Hypervalent iodine reagents became versatile reagents in organic chemistry over the last decades. The mild reaction conditions associated with the low toxicity and the environmentally friendly behaviour of those compounds render them attractive to use in organic synthesis.^[1, 2] Those reagents are very selective oxidants^[3] and several derivatives have been reported as enantiomerically pure reagents.^[4] Due to their electrophilic nature and their excellent leaving-group ability, they can react with a broad range of nucleophiles in reactions such as the oxidation of sulfides to sulfoxides,^[5] the dearomatization of phenols,^[6] the α -arylation^[7] and the α -oxygenation^[8] of carbonyl compounds but also in the functionalization of carbon-carbon double bonds (deoxygenation,^[9] deamination,^[10] oxyamination,^[11] iodoamination,^[12] oxytrifluoromethylation,^[13] or amino-fluorination^[14]). The facile generation of cationic intermediates by hypervalent iodine reagents allows either the direct reaction with a nucleophile or the formation of rearranged products^[15] with ring contraction,^[16] ring expansion,^[17] or aryl migration.^[18] Similar rearrangement have previously been reported with some toxic thallium reagents.^[19] Finally, intensive efforts have been made towards the catalytic use of those hypervalent iodine reagents.^[4b, 20]

We have reported the oxidative rearrangement of aryl-substituted unsaturated carboxylic acids to yield furanones^[21] and described the first stereoselective rearrangement mediated by hypervalent iodine reagent on chalcone derivatives.^[18b] More recently, we developed the stereoselective hypervalent iodine-promoted oxidative rearrangement of 1,1-disubstituted alkenes.^[18a] Haruta et al. reported the oxidative 1,2-aryl migration of alkyl aryl ketones to synthesize 2-aryl propanoates using diacetoxy(iodobenzene) in moderate to good yields as shown in Scheme 1.^[22] 2-Aryl alkanooates are direct precursors of non-steroidal anti-inflammatory drugs (NSAIDs). NSAIDs are sold on



Scheme 1. Rearrangement of aryl alkyl ketones.

[a] F. Malmedy, Prof. Dr. T. Wirth
School of Chemistry, Cardiff University
Park Place, Main Building, Cardiff CF10 3AT (UK)
E-mail: wirth@cf.ac.uk

the market as racemates but usually one of the two enantiomers is less active or even causes side effects.^[23] Therefore, it would be of great interest to synthesize them in enantiomerically pure form. Herein, we describe the development of a stereoselective reaction which allows access to 2-aryl alkanooates in good enantioselectivities.

Results and Discussion

In order to evaluate hypervalent reagents and their reaction conditions for the oxidative rearrangement, the reaction of propiophenone 1 with hypervalent iodine reagents under different reaction conditions in the presence of acidic additives was investigated as shown in Table 1.


As we had previously much success using iodine(III) bistriflates as reagents, we performed the reaction by replacing the sulfuric acid additive with triflic acid and trimethylsilyl triflate. We also relied on the past performance of chiral, hypervalent iodine reagents developed by Ishihara for rearrangement reactions.^[18a]

With (diacetoxyiodo)benzene (Table 1, entries 1 and 2) the reactions proceeded very well and the product 2 was observed with very good conversion and acceptable isolated yield considering the volatility of methyl 2-phenylpropanoate 2. The use of the chiral diester 3 provided initial (low) selectivities of the product 2 with 25% and 10% ee for TMSOTf and TfOH as additives, respectively (Table 1, entries 3 and 4). As hypervalent iodine(III) reagents are usually water sensitive, dry HC(OMe)₃ was used as solvent/reagent. Surprisingly, the reaction did not lead to the formation of the desired product but to complete degradation of starting material (Table 1, entry 5). If 10 equivalents of water or 3 equivalents of methanol were added to the dry solvent, the reaction proceeded as before when performed with normal grade HC(OMe)₃ without an inert atmosphere (Table 1, entries 6 and 7). Other additives such as sulfuric acid or para-toluenesulfonic acid led to lower selectivities (Table 1, entries 8 and 9). Also the use of other solvents together with HC(OMe)₃ did reduce yield and selectivity. Different temperatures and hypervalent iodine reagents were investigated next. When the reaction was performed at !208C using reagent 3, the enantioselectivity increased to 40% ee (Table 1, entry 10). The temperature was further lowered down to !488C, but this led only to an incomplete formation of (1,1-dimethoxypropyl)-benzene (Table 1, entry 11).

Supporting information for this article can be found under <http://dx.doi.org/10.1002/chem.201603022>.

The hypervalent iodine(III) reagent 4 containing amide moieties led to product formation with very good conversion and a higher enantioselectivity of 53 % (Table 1, entry 12). The effect of the Lewis acid in combination with reagent 4 was also investigated. The stereoselectivity was as good using

Table 1. Reaction conditions for the stereoselective rearrangement of propiophenone 1 to methyl-2-phenylpropanoate 2.



| Entry | Additive | Iodine(III) reagent | T [8C] | t [h] | ee [%] | Yield [%] (Conversion) ^[a] |
|-------|-------------------------------------------|-----------------------|--------|-------|--------|-----------------------------------------|
| 1 | 2 equiv. TMSOTf | PhI(OAc) ₂ | 20 | 4 | – | 62 (99) |
| 2 | 2 equiv. p-TsOH·H ₂ O | PhI(OAc) ₂ | 20 | 3 | – | 40 (99) |
| 3 | 2 equiv. TMSOTf | 3 | 0 | 22 | 25 (R) | 22 (84) |
| 4 | 2 equiv. TfOH | 3 | 0 | 16 | 10 (R) | 36 (89) |
| 5 | 2 equiv. TfOH | 3 | 0 | 24 | – | 0 ^[b] |
| 6 | 2 equiv. TfOH, 10 equiv. H ₂ O | 3 | 20 | 24 | 33 (R) | 34 (61) ^[b] |
| 7 | 2 equiv. TfOH, 3 equiv. MeOH | 3 | 20 | 23 | 33 (R) | n.d. ^[c] (66) ^[b] |
| 8 | 2 equiv. H ₂ SO ₄ | 3 | 20 | 23 | 23 (R) | 44 (77) |
| 9 | 2 equiv. p-TsOH·H ₂ O | 3 | 20 | 23 | 13 (R) | 48 (75) |
| 10 | 2 equiv. TfOH | 3 | !20 | 24 | 40 (R) | n.d. ^[c] (96) |
| 11 | 2 equiv. TfOH | 3 | !48 | 10 | – | – ^[d] |
| 12 | 2 equiv. TfOH | 4 | !20 | 24 | 53 (R) | 49 (91) |
| 13 | 2 equiv. TMSOTf | 4 | !20 | 24 | 51 (R) | 18 (39) |
| 14 | 2 equiv. TfOH | 5 | 0 | 23 | 15 (S) | 69(93) |

[a] Conversion determined by ¹H NMR of the crude reaction mixture. [b] Dry HC(OMe)₃ used as solvent. [c] n.d.: not determined. [d] (1,1-Dimethoxypropyl)benzene was isolated as product in 57% yield.

TMSOTf (51% ee) but the conversion dropped to 39% (Table 1, entry 13). When BF₃·OEt₂ was used, no reaction occurred and starting material was recovered. With the new, pyridine-based chiral hypervalent iodine reagent 5^[24] the enantioselectivity was very low (15% ee) and the conversion stayed as high as with the other iodine(III) reagents (Table 1, entry 14) (Figure 1).




Figure 1. Selected chiral hypervalent iodine reagents.

As the volatility of methyl-2-phenylpropanoate gives erroneous results, we investigated the rearrangement of the much less volatile 1-(naphthalen-1-yl)propan-1-one 6, which is easily accessible by a Friedel–Crafts acylation.^[25]


The reaction using reagent 4 worked well and product 7 was isolated in 91% yield with a selectivity of 46% ee when performed under identical reaction conditions to propiophenone. Reagent 3 is less efficient (Table 2, entry 1) and also catalytic amounts of Lewis acid are not enough (Table 2, entry 3) indicating that the Lewis acid must fully activate the iodine(III) reagent and is not only assisting in the equilibrium between the ketone and the enol form.

The influence of the nature of the orthoester was then investigated. Triethyl orthoformate increased the selectivity to 62% ee without drastically affecting the yield (Table 2, entry 4).

This higher enantioselectivity can be explained by the higher bulkiness around the reactive centre where the rearrangement occurs. Therefore, an even bulkier orthoester should induce higher selectivity. However, the reaction with the tri-iso-propyl orthoformate did not lead to any rearranged product. It only added the iso-propoxy group to the α-position of the ketone in 22% yield (Table 2, entry 5). This might be due to the too high bulkiness of the iso-propyl moiety. Indeed, the formation of the ketal is more favoured in the case of the less bulky trimethyl or triethyl orthoformate. If the equilibrium is shifted to the ketal form (R = Me, Et), the presence of two oxygens will favour the rearranged transformation as they can stabilize the positive charge resulting from the aryl group's migration. On the other hand, if the equilibrium is shifted towards the ketone form (R = iPr), the free alcohol is more likely to attack the carbon in α-position to generate the α-alkoxylated product.

Finally, the effect of the temperature on the selectivity was also investigated. At room temperature, the reaction worked similarly well, providing 7 in 81% yield and 44% ee (Table 2, entry 6). At higher temperatures, however, the yield dropped due to side reactions (Table 2, entry 7).

Table 2. Reaction conditions for the stereoselective rearrangement of naphthalene derivative 6 to alkyl 2-(naphthalen-1-yl)propanoate 7.



| Entry | R | Iodine(III) reagent | T [8C] | Product 7 | ee [%] | Yield [%] |
|-------|-----|---------------------|--------|-----------|--------|------------------|
| 1 | Me | 3 | !20 | 7a | 22 (R) | 62 |
| 2 | Me | 4 | !20 | 7a | 46 (R) | 91 |
| 3 | Me | 4 | !20 | 7a | – | 0 ^[a] |
| 4 | Et | 4 | !20 | 7b | 62 (R) | 70 |
| 5 | iPr | 4 | !20 | 7c | – | 0 ^[b] |
| 6 | Me | 4 | 20 | 7a | 44 (R) | 81 |
| 7 | Me | 4 | 40 | 7a | 40 (R) | 40 |

7a (R = Me)
7b (R = Et)
7c (R = i-Pr)

[a] 0.2 equiv. TfOH used. [b] 2-Isopropoxy-1-(naphthalen-1-yl)propan-1-one isolated in 22% yield.

Subsequently, a range of other substrates was investigated and the results are summarized in Table 3. When butyrophe- none was used instead of propiophenone (Table 1, entry 12), the selectivity increased by 10% to reach 63% ee (Table 3, entry 1) and the bulkier isovalerophenone led to a rearranged reaction product with 73% ee (Table 3, entry 2). Unfortunately, as the selectivity increased with the bulkiness of the alkyl chain, the conversion dropped significantly. The reaction with isovalerophenone was also performed at room temperature

Table 3. Stereoselective rearrangements of aryl ketones 8.

| Entry | Ar | R' | R | T | 9 | ee [%] | Yield [%] (racemate) ^[a] |
|-------|--------------------------------------------------|-------------------------------------|----|-----|-----|------------------|----------------------------------------|
| | | | | | | [8C] | |
| 1 | Ph | Et | Me | !20 | 9a | 63 (R) | 56 (58) |
| 2 | Ph | iPr | Me | !20 | 9b | 73 (R) | 31 |
| 3 | Ph | iPr | Me | 20 | 9b | 68 (R) | 31 (93) |
| 4 | Ph | iPr | Et | 20 | 9bb | 35 (R) | 31 (80) |
| 5 | 4-Me-C ₆ H ₄ | Me | Me | 20 | 9c | 46 | 79 (86) |
| 6 | 4-Et-C ₆ H ₄ | Me | Me | 20 | 9d | 44 | 87 (91) |
| 7 | 4-iBu-C ₆ H ₄ | Me | Me | 20 | 9e | 35 | 72 (94) |
| 8 | 4-MeO-C ₆ H ₄ | Me | Me | 20 | 9f | 27 | 80 (82) |
| 9 | 1-(6-MeO-naphth) | Me | Me | 20 | 9g | – | (traces) |
| 10 | 4-F-C ₆ H ₄ | Me | Me | !20 | 9h | 45 | 50 (92) |
| 11 | 2-Br-C ₆ H ₄ | Me | Me | 20 | 9i | 44 | 46 (79) |
| 12 | 3-Br-C ₆ H ₄ | Me | Me | 20 | 9j | 46 | 14 (98) |
| 13 | 4-Br-C ₆ H ₄ | Me | Me | 20 | 9k | – ^[b] | 28 (76) |
| 14 | 3-CF ₃ -C ₆ H ₄ | Me | Et | 20 | 9l | – | 13 (23) ^[c] |
| 15 | 3-NO ₂ -C ₆ H ₄ | Me | Me | 20 | 9m | – | n.r. ^[d] (75) |
| 16 | 2-pyridyl | Me | Me | 20 | 9n | – | – ^[e] |
| 17 | Ph | cyclo-C ₃ H ₅ | Et | 20 | 9o | – | – ^[f] |
| 18 | 2-naphthyl | Me | Me | !20 | 9p | 24 | 52 (83) ^[g] |

[a] Yield for the racemate using PhI(OAc)₂ as reagent. [b] ee could not be determined by HPLC as the enantiomers could not be separated. The product showed optical rotation. [c] 2-Ethoxy-1-(3-(trifluoromethyl)phenyl)propan-1-one was isolated as only product. With PhI(OAc)₂ also 73% rearranged product are obtained. [d] n.r.: no reaction occurred with reagent 4. [e] 2-(1,1-Dimethoxypropyl)pyridine trifluoromethanesulfonic acid adduct was isolated as only product. [f] 4-Ethoxy-1-phenylbutan-1-one as ring-opened product was isolated in 98% yield. Further rearrangement occurred only with PhI(OAc)₂, not with 4. [g] Reagent 3 was used instead of 4.


whereby the enantioselectivity dropped only by 5% and confirmed the small impact of the temperature on the rearrangement (Table 3, entry 3). The reaction was also carried out with triethyl orthoformate as it was beneficial for the rearrangement of 1-propionynaphthalene. However, with isovalerophenone the enantioselectivity dropped to 35% ee (Table 3, entry 4).

Subsequently, the influence of the electronic properties of the aromatic ring in the rearrangement process was investigated. Propiophenone derivatives with electron-rich aryl moieties gave the rearranged products in very good yields but only moderate enantioselectivities (Table 3, entries 5–7). Hydrolysis of the methyl ester product shown in Table 3, entry 7, was achieved with 1N sodium hydroxide in THF/methanol to yield ibuprofen in 93% yield, but with reduced enantioselectivity (22% ee). The reaction proceeded similarly well with 4-methoxypropionophenone, but surprisingly did not work with 1-(6-methoxynaphthalen-1-yl)propan-1-one (Table 3, entries 8 and 9).

In the reaction with 4-methoxypropionophenone, the reaction product 9f reacted further with an excess of (diacetoxyiodo)benzene to generate the diaryliodonium salt 9ff, its structure was also confirmed by X-ray analysis^[24] (see Scheme 2 and the Supporting Information). Compound 9f can also be converted into 9ff under the reaction conditions in 90% yield. The formation of 9ff can be suppressed completely when only one equivalent of (diacetoxyiodo)benzene is used.

Electron-withdrawing groups on the aromatic moiety do not seem to affect the enantioselectivity. Indeed, for all the compounds for which the enantioselectivity could be determined, the enantiomeric excess was between 44 and 46% (Table 3, entries 10–12). However, the yield of the reaction is depending on the nature of the electron-withdrawing group. Compared to fluorine, the yield decreased to 28% when a bromine substituent is in para-position (Table 3, entries 10 and 13). The position on the aromatic ring is also important. Whereas para- and meta-substituents gave poor yields (28 and 14%, respectively), the ortho-substituted product can be obtained in moderate yield (46%) (Table 3, entries 11–13). With a trifluoromethyl substituent in the meta-position, the rearranged product could only be obtained in trace amounts and the α -ethoxylated product was obtained as the major product in 13% yield (Table 3, entry 14). 3-Nitropropionophenone was not reactive enough (Table 3, entry 15) and a pyridine derivative led only to the corresponding ketal (Table 3, entry 16). A cyclopropyl-substituted derivative ring-opened but did not rearrange with the chiral reagent 4 (Table 3, entry 17).


Electron-rich aryl moieties migrate faster as they stabilize the intermediate phenonium ion. However, the enantioselectivity seems to follow the opposite trend. This may be due to the fact that the reaction rate is faster for electron-rich aryl moieties and the interaction with the chiral reagent is less strong to induce high selectivity.




Scheme 2. Rearrangement and formation of diaryliodonium salt 9ff.

The proposed mechanism for the rearrangement as shown in Scheme 3 is similar to the mechanism proposed by Haruta.^[22] In the generation of intermediate 11 the stereochemistry of the rearrangement process is defined. The presence of large amounts of nucleophiles will directly interfere with the intermediate 11 in a direct substitution of the iodine(III) moiety leading to α -functionalized ketones, as we and others have shown previously. Protected enolethers have been used by us and others recently as efficient substrates for stereoselective iodine(III)-mediated reactions. Under the reaction conditions employed, it is also plausible that the presence of triflic acid and trimethyl orthoformate will lead to the formation of the enol 15a via ketal 14.^[26] Although it is not identifiable in the NMR,^[8a] it could react with the iodine(III) species in the presence of methanol to form intermediate 12.

As shown in Scheme 4, silylenolether 15b provides a mixture of rearranged product 2 and direct substitution product 16,




Scheme 3. Proposed mechanism for the rearrangement.



Scheme 4. Enols and ketals as substrates for the rearrangement reaction.

whereas the acetyl-substituted enolether 15 c ($R = \text{Ac}$) only forms α -substituted product 16 (57% yield using $\text{PhI}(\text{OAc})_2$), probably because the formation of the corresponding ketal is less efficient. The pre-generated ketal 14 is similarly efficient in the rearrangement reaction and gives the rearrangement product 2 in similar conversion and selectivity to propiophenone. A rapid interconversion between 1 and 14 via the hemiketal under the reaction conditions was confirmed by NMR spectroscopy (Supporting Information).

The absolute stereochemistry of the products 2 and 16 indicates the common intermediate 12 resulting from a Si-attack of the iodine(III) electrophile 4 to the enol 10/15 a. Reaction by rearrangement (path A) generates compound 2 with (*R*)-configuration and backside substitution with methanol (path B) leads to 16 with (*S*)-configuration. The different stereochemical descriptors are due to the Cahn–Ingold–Prelog priorities and not to different stereochemical pathways. The absolute configuration of 16 was confirmed by its independent synthesis from lactic acid (Supporting Information). (Figure 2)


Figure 2. Intermediate 12 ($R =$

H, Me).

Other substrates, however, seem not to be suitable for the rearrangement reaction. Compounds

of type 17 with substituents R changing the electronic properties only lead to the α -ethoxylated

products (Scheme 5). Only in the case of an aryl substitution such as in 19 a the rearranged product 20 a is formed together



Scheme 5. Substituted substrates for the rearrangement reaction.

ranged compound (21 a) and some α -methoxylated product 22 a. The selectivities obtained with reagent 4 are 25 % ee (62 % yield) for 20 a, 5 % ee (4 % yield) for 21 a and 7 % ee (11 % yield) for 22 a, but the absolute configurations of the products could not be determined. Similar yields (20 a: 70 %, 21 a: 10 %, 22 a: 10 %) are obtained when (diacetoxyiodo)benzene is employed as iodine(III) reagent. Compound 19 b ($R'' = \text{Me}$), which would lead to products with stereogenic tetrasubstituted carbon atoms, is unreactive and completely recovered under the reaction conditions.

The catalytic transformation was investigated using $\text{PhI}(\text{OAc})_2$ as the in situ synthesized hypervalent iodine reagent. Different oxidants ($\text{NaBO}_3 \cdot 4\text{H}_2\text{O}$, *m*-CPBA) were investigated together with iodobenzene, propiophenone and trimethyl orthoformate, but no rearranged product was obtained under the reaction conditions investigated.

Conclusion

In summary, we have developed a stereoselective rearrangement of aryl alkyl ketones mediated by chiral hypervalent iodine (III) reagents. 2-Arylpropionate derivatives were synthesized in moderate to good yields with enantioselectivities up to 73 % using a lactic acid-based reagent in the presence of TfOH and trimethyl orthoformate. Further investigations to improve the enantioselectivity and the development of a catalytic protocol are in progress.

Experimental Section

Rearrangement of propiophenone derivatives: To a solution of propiophenone derivative 8 (0.305 mmol) and the Ishihara amide 4 (267 mg, 0.365 mmol) in $\text{HC}(\text{OMe})_3$ (1.5 mL), TfOH (57 mL, 0.609 mmol) was added dropwise at the temperature mentioned in Table 3. The resulting mixture was stirred at this temperature for 48 h, quenched with water (1 mL), extracted with CH_2Cl_2 (3 \times 2 mL), dried over a Telos phase separator and the solvent was removed under reduced pressure. After column chromatography (0 to 10 % Et_2O in hexane), the corresponding product 9 was obtained.

Acknowledgements

with methoxylation at the phenyl substituent of the rear-

We thank Dr. B. Kariuki, Cardiff University, for the X-ray analysis of compounds 5 and 9 ff. Support from the School of Chemis-

Full Paper


try, Cardiff University, is gratefully acknowledged. We thank the EPSRC National Mass Spectrometry Facility, Swansea, for mass spectrometric data.

Keywords : addition · alkenes · iodine(III) · rearrangement · stereoselective synthesis

- [1] a) T. Wirth, *Hypervalent Iodine Chemistry*, Vol. 224 (Top. Curr. Chem.), Springer, Berlin, 2003; b) T. Wirth, *Hypervalent Iodine Chemistry*, Vol. 373 (Top. Curr. Chem.), Springer, Berlin, 2016.
- [2] V. V. Zhdankin, *Hypervalent Iodine Chemistry*, Wiley, Chichester, 2014.
- [3] T. Wirth, *Angew. Chem. Int. Ed.* 2005, 44, 3656–3665; *Angew. Chem.* 2005, 117, 3722–3731.
- [4] a) R. Kumar, T. Wirth, *Top. Curr. Chem.* 2015, 373, 243–261; b) F. Berthiol, *Synthesis* 2015, 587–603; c) F. V. Singh, T. Wirth, *Chem. Asian J.* 2014, 9, 950–971; d) R. M. Romero, T. H. Wçste, K. Muçiz, *Chem. Asian J.* 2014, 9, 972–983; e) A. Parra, S. Reboredo, *Chem. Eur. J.* 2013, 19, 17244–17260; f) U. Farid, T. Wirth in *Asymmetric Synthesis – The Essentials* (Eds.: S. Br#se, C. Christmann), Wiley-VCH, 2012, 197–203; g) M. Uyanik, K. Ishihara, *J. Synth. Org. Chem. Jpn.* 2012, 70, 1116–1122; h) H. Liang, M. A. Ciuffolini, *Angew. Chem. Int. Ed.* 2011, 50, 11849–11851; *Angew. Chem.* 2011, 123, 12051–12053; i) M. Ngatimin, D. W. Lupton, *Aust. J. Chem.* 2010, 63, 653–658.
- [5] a) S. M. Altermann, S. Sch#fer, T. Wirth, *Tetrahedron* 2010, 66, 5902–5907; b) V. V. Zhdankin, J. T. Smart, P. Zhao, P. Kiprof, *Tetrahedron Lett.* 2000, 41, 5299–5302; c) H. Tohma, S. Takizawa, H. Watanabe, Y. Fukuoaka, T. Maegawa, Y. Kita, *J. Org. Chem.* 1999, 64, 3519–3523; d) D. G. Ray, G. F. Koser, *J. Am. Chem. Soc.* 1990, 112, 5672–5673; e) T. Imamoto, H. Koto, *Chem. Lett.* 1986, 967–968.
- [6] a) A. M. Harned, *Tetrahedron Lett.* 2014, 55, 4681–4689; b) K. A. Volp, A. M. Harned, *Chem. Commun.* 2013, 49, 3001–3003; c) M. Uyanik, T. Yasui, K. Ishihara, *Angew. Chem. Int. Ed.* 2013, 52, 9215–9218; *Angew. Chem.* 2013, 125, 9385–9388; d) T. Dohi, N. Takenaga, T. Nakae, Y. Toyoda, M. Yamasaki, M. Shiro, H. Fujioka, A. Maruyama, Y. Kita, *J. Am. Chem. Soc.* 2013, 135, 4558–4566; e) S. Quideau, G. Lyvinec, M. Marguerit, K. Bathany, A. Ozanne-Beaudenon, T. Buffeteau, D. Cavagnat, A. Ch#ned\$, *Angew. Chem. Int. Ed.* 2009, 48, 4605–4609; *Angew. Chem.* 2009, 121, 4675–4679.
- [7] a) M. Brown, M. Delorme, F. Malmedy, J. Malmgren, B. Olofsson, T. Wirth, *Synlett* 2015, 1573–1577; b) V. K. Aggarwal, B. Olofsson, *Angew. Chem. Int. Ed.* 2005, 44, 5516–5519; *Angew. Chem.* 2005, 117, 5652–5655; c) M. Ochiai, Y. Kitagawa, N. Takayama, Y. Takaoka, M. Shiro, *J. Am. Chem. Soc.* 1999, 121, 9233–9234.
- [8] a) B. Basdevant, C. Y. Legault, *J. Org. Chem.* 2015, 80, 6897–6902; b) B. Basdevant, C. Y. Legault, *Org. Lett.* 2015, 17, 4918–4921; c) P. Mizar, T. Wirth, *Angew. Chem. Int. Ed.* 2014, 53, 5993–5997; *Angew. Chem.* 2014, 126, 6103–6107; d) U. Farooq, S. Sch#fer, A.-H. Shah, D. Freudendahl, T. Wirth, *Synthesis* 2010, 1023–1029; e) S. M. Altermann, R. D. Richardson, T. K. Page, R. K. Schmidt, E. Holland, U. Mohammed, S. M. Paradine, A. N. French, C. Richter, A. M. Bahar, B. Witulski, T. Wirth, *Eur. J. Org. Chem.* 2008, 5315–5328; f) R. Richardson, T. Page, S. Altermann, S. Paradine, A. French, T. Wirth, *Synlett* 2007, 538–542; E. Hatzigrigoriou, A. Varvoglis, M. Bakola-Christianopoulou, *J. Org. Chem.* 1990, 55, 315–318.
- [9] a) S. Haubenreisser, T. H. Wçste, C. Mart#nez, K. Ishihara, K. Muçiz, *Angew. Chem. Int. Ed.* 2016, 55, 413–417; *Angew. Chem.* 2016, 128, 422–426; b) T. Takesue, M. Fujita, T. Sugimura, H. Akutsu, *Org. Lett.* 2014, 16, 4634–4637; c) M. Shimogaki, M. Fujita, T. Sugimura, *Eur. J. Org. Chem.* 2013, 7128–7138; d) M. Fujita, K. Mori, M. Shimogaki, T. Sugimura, *Org. Lett.* 2012, 14, 1294–1297; e) M. Fujita, Y. Yoshida, K. Miyata, A. Wakisaka, T. Sugimura, *Angew. Chem. Int. Ed.* 2010, 49, 7068–7071; *Angew. Chem.* 2010, 122, 7222–7225; f) M. Fujita, Y. Ookubo, T. Sugimura, *Tetrahedron Lett.* 2009, 50, 1298–1300; g) M. Fujita, S. Okuno, H. J. Lee, T. Sugimura, T. Okuyama, *Tetrahedron Lett.* 2007, 48, 8691–8694; h) U. H. Hirt, M. F. H. Schuster, A. N. French, O. G. Wiest, T. Wirth, *Eur. J. Org. Chem.* 2001, 1569–1579; i) T. Wirth, U. H. Hirt, *Tetrahedron: Asymmetry* 1997, 8, 23–26.
- [10] a) P. Mizar, A. Laverny, M. El-Sherbini, U. Farid, M. Brown, F. Malmedy, T. Wirth, *Chem. Eur. J.* 2014, 20, 9910–9913; b) C. Rçben, J. A. Souto, E. C. Escudero-Adán, K. Muçiz, *Org. Lett.* 2013, 15, 1008–1011; c) C. Rçben, J. A. Souto, Y. González, A. Lishchynskiy, K. Muçiz, *Angew. Chem. Int. Ed.* 2011, 50, 9478–9482; *Angew. Chem.* 2011, 123, 9650–9654.
- [11] U. Farid, T. Wirth, *Angew. Chem. Int. Ed.* 2012, 51, 3462–3465; *Angew. Chem.* 2012, 124, 3518–3522.
- [12] P. Mizar, A. Burrelli, E. G'nther, M. Sçftje, U. Farooq, T. Wirth, *Chem. Eur. J.* 2014, 20, 13113–13116.
- [13] R. Zhu, S. L. Buchwald, *Angew. Chem. Int. Ed.* 2013, 52, 12655–12658; *Angew. Chem.* 2013, 125, 12887–12890.
- [14] a) H. Chen, A. Kaga, S. Chiba, *Org. Biomol. Chem.* 2016, 14, 5481–5485; b) W. Kong, P. Feige, T. de Haro, C. Nevado, *Angew. Chem. Int. Ed.* 2013, 52, 2469–2473; *Angew. Chem.* 2013, 125, 2529–2533.
- [15] a) G. Maertens, S. Canesi, *Top. Curr. Chem.* 2015, 373, 223–242; b) S. Desjardins, G. Maertens, S. Canesi, *Org. Lett.* 2014, 16, 4928–4931; c) F. V. Singh, T. Wirth, *Synthesis* 2013, 2499–2511; d) G. Jacquemot, S. Canesi, *J. Org. Chem.* 2012, 77, 7588–7594; e) M.-A. Beaulieu, K. C. Gu#rdard, G. Maertens, C. Sabot, S. Canesi, *J. Org. Chem.* 2011, 76, 9460–9471; f) M.-A. Beaulieu, C. Sabot, N. Achache, K. C. Gu#rdard, S. Canesi, *Chem. Eur. J.* 2010, 16, 11224–11228; g) A. A. Zagulyaeva, C. T. Banek, M. S. Yusubov, V. V. Zhdankin, *Org. Lett.* 2010, 12, 4644–4647; h) J.-M. Vat(, *Synlett* 2008, 1785–1788.
- [16] a) L. F. Silva, F. A. Siqueira, E. C. Pedrozo, F. Y. M. Vieira, A. C. Doriguetto, *Org. Lett.* 2007, 9, 1433–1436; b) L. F. Silva, *Molecules* 2006, 11, 421–434; c) F. A. Siqueira, E. E. Ishikawa, A. Fogaãa, A. T. Faccio, V. M. T. Carneiro, R. R. S. Soares, A. Utaka, I. R. M. T#b\$ka, M. Bielawski, B. Olofsson, L. F. Silva, *J. Braz. Chem. Soc.* 2011, 22, 1795–1807.
- [17] a) L. F. Silva, R. S. Vasconcelos, M. A. Nogueira, *Org. Lett.* 2008, 10, 1017–1020; b) M. Justik, G. Koser, *Molecules* 2005, 10, 217–225.
- [18] a) M. Brown, R. Kumar, J. Rehbein, T. Wirth, *Chem. Eur. J.* 2016, 22, 4030–4035; b) U. Farid, F. Malmedy, R. Claveau, L. Albers, T. Wirth, *Angew. Chem. Int. Ed.* 2013, 52, 7018–7022; *Angew. Chem.* 2013, 125, 7156–7160; c) A. C. Boye, D. Meyer, C. K. Ingison, A. N. French, T. Wirth, *Org. Lett.* 2003, 5, 2157–2159.
- [19] a) H. M. Ferraz, A. M. Aguilar, L. F. Silva, *Synthesis* 2003, 1031–1034; b) A. McKillop, B. P. Swann, E. C. Taylor, *J. Am. Chem. Soc.* 1973, 95, 3340–3343.
- [20] R. D. Richardson, T. Wirth, *Angew. Chem. Int. Ed.* 2006, 45, 4402–4404; *Angew. Chem.* 2006, 118, 4510–4512.
- [21] F. V. Singh, J. Rehbein, T. Wirth, *ChemistryOpen* 2012, 1, 245–250.
- [22] Y. Tamura, T. Yakura, Y. Shirouchi, J.-I. Haruta, *Chem. Pharm. Bull.* 1985, 33, 1097–1103.
- [23] a) S. S. Adams, P. Bresloff, C. G. Mason, *Curr. Med. Res. Opin.* 2008, 3, 552–552; b) I. Agranat, H. Caner, J. Caldwell, *Nat. Rev. Drug Discovery* 2002, 1, 753–768; c) A. H. Beckett, *Biochem. Soc. Trans.* 1991, 19, 443–446.
- [24] CCDC 1487361 and 1487362 for 5 and 9 ff, respectively, contain the supplementary crystallographic data for this paper. These data are provided free of charge by The Cambridge Crystallographic Data Centre
- [25] A. K. Ghosh, J. Takayama, Y. Aubin, K. Ratia, R. Chaudhuri, Y. Baez, K. Sleeman, M. Coughlin, D. B. Nichols, D. C. Mulhearn, B. S. Prabhakar, S. C. Baker, M. E. Johnson, A. D. Mesecar, *J. Med. Chem.* 2009, 52, 5228–5240.
- [26] a) S. Manojveer, R. Balamurugan, *Org. Lett.* 2014, 16, 1712–1715; b) F. Wu, R. Bai, Y. Gu, *Adv. Synth. Catal.* 2016, 358, 2307–2316.

Received: June 24, 2016
Published online on && &&, 0000

FULL PAPER



& Stereoselective Synthesis


F. Malmedy, T. Wirth*

&& - &&

Stereoselective Ketone
Rearrangements with Hypervalent
Iodine Reagents

Now stereoselective: Iodine(III)-mediated rearrangement of arylketones in the presence of orthoesters are possible

leading to α -arylated esters in selectivities of up to 73% ee.



Stereoselective Rearrangements

The first stereoselective version of an iodine(III)-mediated rearrangement of arylketones in the presence of orthoesters is described T. Wirth and F. Malmedy on page && ff. The reaction products, α -arylated esters, are very useful intermediates in the synthesis of bioactive compounds such as ibuprofen. With chiral lactic acid-based iodine(III) reagents, enantioselectivities of up to 73% ee have been achieved.