

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/96548/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Parry, Elinor, Gregory, S. D., Lauridsen, R. B. and Griffiths, Sian 2018. The effects of flow on Atlantic salmon (*Salmo salar*) redd distribution in a UK chalk stream between 1980 and 2015. *Ecology of Freshwater Fish* 27 (1) , pp. 128-137. 10.1111/eff.12330

Publishers page: <http://dx.doi.org/10.1111/eff.12330>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





The effects of flow on Atlantic salmon (*Salmo salar*) redd distribution in a UK chalk stream between 1980 and 2015

Journal:	<i>Ecology of Freshwater Fish</i>
Manuscript ID	EFF-16-0075.R2
Manuscript Type:	Original articles
Date Submitted by the Author:	17-Nov-2016
Complete List of Authors:	Parry, Elinor; Game and Wildlife Conservation Trust, Salmon and Trout Research Centre; Cardiff University Cardiff School of Biosciences, Cardiff School of Biosciences Gregory, Stephen; Game and Wildlife Conservation Trust, Salmon and Trout Research Centre Lauridsen, Rasmus; Game and Wildlife Conservation Trust, Salmon and Trout Research Centre Griffiths, Sian; Cardiff University Cardiff School of Biosciences
Keywords:	Atlantic salmon, Redd distribution, Climate change, Abstraction, River discharge

SCHOLARONE™
Manuscripts

Preprint Only

1 The effects of flow on Atlantic salmon (*Salmo salar*) redd
2 distribution in a UK chalk stream between 1980 and 2015

3
4 Elinor S. Parry^{1,2}, Stephen D. Gregory¹, Rasmus B. Lauridsen¹, Siân W. Griffiths²

5 1 Game and Wildlife Conservation Trust, Salmon and Trout Research Centre, Wareham, Dorset BH20
6 6BB, United Kingdom.

7 2 Cardiff School of Biosciences, Cardiff University, Sir Martin Evans Building, Museum Avenue, Cardiff
8 CF10 3AX, United Kingdom.

9 Corresponding author: esparry@msn.com, telephone +441597824050

10 Coauthor emails: Stephen Gregory (sgregory@gwct.org.uk), Rasmus Lauridsen
11 (rlauridsen@gwct.org.uk), Siân Griffiths (GriffithsSW@cardiff.ac.uk)

12 Running head: Flow affects salmon redd distribution

13 Keywords: Atlantic salmon, spawning migration, river discharge, redd distribution, climate change,
14 abstraction

15 Word counts (excluding headings and equations):

16 Abstract = 249

17 Main text = 4249 (Introduction = 1004; Methods = 1265; Results = 648; Discussion &
18 Conclusion = 1877)

19 Figures and Tables 7 (+ 10 Supporting Information)

20 References = 53

21

22 Abstract

23 Atlantic salmon are an ecologically and economically important migratory fish in the UK, whose
24 stocks have been declining over the past 30 years. Future climate and water use changes have the
25 potential to alter the reproductive behaviour and distribution of salmon within a river, by restricting
26 times and ability to access suitable spawning areas. As the survival of emergent salmon juveniles is
27 density dependent, understanding how climate-driven changes in flow affect the location of salmon
28 redds is important for future conservation efforts. This study examined how flow conditions affect
29 the distribution of redds within a UK chalk stream, the river Frome in Dorset. Sixteen years of redd
30 distribution and flow data between 1980 and 2015 were analysed using Linear Mixed Effects
31 modelling. Generally, highest redd densities occurred within middle reaches of the main river. Mean
32 flow during the river Frome critical migration period (October – December) did not affect the density
33 of redds directly but affected the relationship between redd density and distance from tidal limit:
34 redd densities were spread more uniformly throughout the river under high flow conditions whereas
35 redds were more aggregated in the middle river reaches under low flow conditions. Together, these
36 findings suggest that access to upstream spawning grounds was limited under low flow conditions,
37 which could have negative repercussions on juvenile survival. This study has revealed the
38 distribution of redds along the river Frome for the first time and provided a basis for further study
39 into the effects of redd distribution on subsequent juvenile life-stages.

40

41 Introduction

42 Migration between feeding grounds and suitable spawning areas is an important event in the life
43 history of several fish species, having profound survival and fitness implications (Gross et al., 1988).
44 For some species, such as the semelparous European eel (*Anguilla anguilla*), migration to spawning
45 grounds is their only opportunity to contribute to the next generation. For other long distance
46 migrants, such as the Atlantic salmon (*Salmo salar*; hereafter termed salmon), multi-annual
47 spawning is possible and reproductive success is in part dependent on other factors, such as
48 competition for spawning habitat and mates (Fleming, 1998). During salmonid migration,
49 movements made to find habitat suitable for nest or redd construction is known as the search phase
50 of migration (*sensu* Thorstad et al. 2008). Although several studies have investigated the influence of
51 environmental conditions on the timing and success of migratory behaviour in the period preceding
52 the search phase, (known as the upstream phase, see reviews by Thorstad et al. 2008; Jonsson &
53 Jonsson 2009; Warren et al. 2015), fewer studies have investigated the role played by environmental
54 conditions on the final distribution of redds throughout rivers, i.e. the outcome of the search phase
55 (Chapman et al., 1986, Klett et al., 2013). This is somewhat surprising given the importance of the
56 search phase for successful reproduction and fitness.

57 Studies to date have revealed that salmon construct redds in fine river gravels at the upstream part
58 of riffles, which provide optimum oxygenation for incubating their eggs and embryos (Bardonnnet and
59 Baglinière 2000). Spawning close to deep pools or vegetative cover can also be advantageous as
60 these habitats can reduce predation of breeding adults and developing juveniles (White, 1942 as
61 cited in Armstrong et al., 2003). Other environmental factors, including temperature, sedimentation,
62 river depth and water velocity can also affect the location of a salmon redd (Moir et al., 1998, Moir
63 et al., 2002, Jonsson and Jonsson, 2009, Armstrong et al., 2003). For example, water velocity and
64 depth, as well as gravel size, are important factors for Chinook salmon (*Oncorhynchus tshawytscha*)
65 redd site selection (Groves and Chandler 1999; Kondolf and Wolman 1993).

66 Climate change is considered one of the greatest threats to biodiversity in the 21st century (Bellard
67 et al., 2012, Thuiller, 2007). Rivers are less stable than marine ecosystems and therefore more prone
68 to rapid changes in conditions, such as temperature or flow (Armstrong et al., 2003). Under current
69 climate change predictions, the number of extreme weather events will increase over the next
70 century, which may directly impact rivers through increases in extreme flow and temperature events
71 (Hulme et al., 2002). Climate driven changes in extreme flow might affect salmon spawning success
72 in a number of ways. Extreme high flow can result in gravel washout and damage to existing redds,
73 while sedimentation of redds following high flows can negatively affect egg and embryo
74 development and subsequent juvenile survival (Levasseur et al., 2006, Lapointe et al., 2000). On the
75 other hand, at extreme low flow, factors such as reduced oxygen content and increased temperature
76 may also hinder upstream migration (Solomon and Sandbrook, 2004). At the very extreme, drought
77 conditions could result in the drying out of redds and subsequent mortality of developing embryos
78 (Becker and Neitzel, 1985).

79 While the flow requirements and other preferences of spawning salmon have been studied
80 extensively, little research has focused on the impact of flow on the overall distribution and density
81 of redds along a river. It has been suggested that reduced flow during the period of peak migration
82 can slow the upstream phase of migration, particularly of larger adults that preferentially spawn
83 further upstream, leading to spawning aggregations below within-river barriers (Mitchell, et al. 2007;
84 Solomon & Sambrook 1999). Studies on the river Tay and Dee in Scotland suggest that increased
85 water discharge during the period of final migration, here assumed to be the search phase of
86 migration, allowed salmon to access shallower spawning tributaries that did not provide suitable
87 conditions for residence earlier in the season (Hawkins 1989; Webb 1989; Webb and Hawkins 1989).
88 Aside from these studies, there are few studies that have directly investigated the effects of flows –
89 and particularly extreme flows – on annual salmon redd distributions. Geist & Dauble (1998)
90 examined the role of river geomorphology and its interaction with microhabitat variables in
91 explaining Chinook salmon redd distributions, but their contrasts were different rivers. Chapman et

92 al. (1986) examined the influence of flow on annual variation in the morphology and density of
93 Chinook salmon redds on the Columbia river but their study was done at a single location. The
94 findings from these studies should arguably be compared with caution, as the species and riverine
95 hydrology can vary significantly compared to British or European rivers.

96 Using one of the longest running datasets on salmonid redds in the UK, we investigate how river
97 flow affects the distribution of salmon redds on a British chalk stream. We build and compare
98 statistical models to describe observed inter-annual changes in salmon redd densities at multiple
99 survey locations on the river Frome, UK, from simple explanatory variables including river flow.
100 Initial analysis of redds along the river Frome suggest that distribution is greatest within the middle
101 reaches, due to the higher abundance of suitable flow conditions and habitat. We predict that low
102 flows during the upstream and search migration phases will result in a higher redd density in the
103 middle relative to the upper sections of the river (Fig. 1). We also predict that as flow increases, redd
104 density will become more uniform across the river (Fig. 1). To our knowledge, this is the first
105 example of how flow conditions during the search phase of Atlantic salmon spawning migration may
106 directly affect the distribution of salmon redds along a British river. Our findings are important due
107 to the density dependent survival of juvenile salmon, which may be governed by the location of
108 redds along the river (Armstrong and Nislow, 2006). Our results may therefore have implications for
109 the future recruitment and survival of salmon within UK rivers under current climate change
110 predictions.

111

112 **[Figure 1 here]**

113

114 Methods

115 Site description and data collection

116 The river Frome, Dorset, UK, is one of 224 British chalk streams, comprising over 80km of braided
117 channels, and is fed primarily by groundwater upwellings from the surrounding chalk and clay
118 geology (Brunner et al., 2010, Sear et al., 1999). The river discharge is therefore alkaline, relatively
119 stable and seldom prone to spates or flooding (National Rivers Authority, 1995). The river Frome has
120 seen considerable anthropomorphic change over the past two centuries, including additions and
121 removals of a number of weirs, fish passes and other modifications to river levels and discharge,
122 such as abstraction (Solomon, 2000).

123 Salmon have been studied for over 50 years on the river Frome (Lauridsen et al., 2015). They
124 reproduce in freshwater between November and February each year. Females excavate a redd in
125 suitable gravel riverbed substrate, into which they deposit their eggs. A description of river Frome
126 salmon redd structure and function (compared with those in other rivers) is given in Crisp & Carling
127 (1989). Male(s) then fertilise the eggs after which the female buries them. Algae and aquatic
128 macrophytes grow slowly in the cold winter water temperature in the river Frome so that redds can
129 be identified from the river bank by their size and presence of clean (i.e., recently disturbed) gravel
130 (Riedl & Peter 2013).

131 Redd surveys

132 River Frome salmonid redd surveys (hereafter *redd surveys*) were started by river wardens in the
133 1950's to monitor salmonid stocks. Initially, these surveys were irregular and inconsistent. Since the
134 1980's, however, the Centre for Ecology and Hydrology and the Environment Agency, with the
135 support from the river Frome, Piddle and West Dorset Fishing Association and the Weld Estate, have
136 carried out comparable salmonid redd surveys throughout the river Frome, from headwaters near
137 Cattistock (section 14 in Fig. 2) downstream to Worgret (section 0 in Fig. 2).

138

139 **[Figure 2 here]**

140

141 Redd surveys were carried out in January or February, dependent on conditions. Two surveyors,
142 equipped with polarised lenses, counted the number of redds within each survey section and
143 estimated their length, width and the cleanliness of riverbed gravel, an indication of the age of the
144 redd. The location of each salmonid redd was recorded; approximate locations were recorded on
145 maps from 1980 to 1992, while the actual location of each redd was recorded on a handheld Global
146 Positioning System for surveys after 2000. Maps indicating approximate redd locations were
147 digitised and georeferenced, allowing the longitude and latitude of each redd to be extracted. Up to
148 a total of 29 survey sections were surveyed in any one year (Fig. 2), although the proportion of
149 sections surveyed in each year varied from 20-93% (mean 53%). Redds more than 110cm wide were
150 deemed to be salmon redds, whereas redds less than 110cm wide were noted as trout (Wessex
151 Water Authority, 1987). Trout redds were omitted from analysis and redd density was calculated as
152 the number of salmon redds divided by the length of the survey section.

153 We calculated two types of explanatory variables for our analysis. Distance from tidal limit (in
154 kilometres) of survey section start and end points were calculated by hand using QGIS software
155 (www.qgis.org) and later verified by the online mapping system MAgiC (magic.defra.gov.uk). The UK
156 Environment Agency has recorded flow at East Stoke on the river Frome every 15 mins since 1965
157 (ES on Fig. 2; see nrfa.ceh.ac.uk/data/station/info/44001). Flow measured at East Stoke was highly
158 correlated with flow measured at Louds Mill, a flow gauge station located approximately 20km
159 upstream (LM on Fig. 2; Pearson's $r = 0.75$, Confidence Intervals: 0.74 - 0.77), confirming that flow
160 measured at East Stoke is representative of flow elsewhere in the catchment. Two flow variables
161 were calculated for this analysis, chosen to represent the critical flow conditions available for
162 spawning migration, shown to be an important determinant of migration success (e.g., Thorstad et

163 al., 2008; Warren et al., 2015): (1) *mean flows* – average flow per year based on mean daily flow
164 between 1st October to 31st December each year, and (2) *high flows* – the number of days between
165 1st October and 31st December that the daily flow exceeded the Q75 flow calculated for the same
166 period in each year. The precise details of these variables, e.g., the dates, were agreed after
167 extensive consultation with local experts about the patterns of spawning salmon migration (Fig. S1).
168 These two flow variables were highly correlated (Pearson's $r = 0.923$) and were thus explored in
169 separate models analysing the effect of flow on redd density.

170 Data analysis

171 A total of 16 years of redd surveys between 1980 and 2014 were chosen for analysis (see Table 1,
172 Appendix). These 16 years were taken from a larger dataset of 22 years between 1957 and 2014
173 because they were surveyed using comparable methods and had the most complete redd survey
174 and river discharge records. Plots of these redd density against distance from tidal limit are
175 presented for these years in Supporting Information (Fig. S2). Of these 16 representative years, five
176 years (1983, 1985, 2007, 2008, 2012) were both high flow years and years in which less than 50% of
177 the sections were surveyed. To test whether inclusion of these years biased our results, we repeated
178 our analysis excluding them and report the results in the Supporting Information. Not all sections
179 were surveyed in all years and the proportion of survey sections surveyed varied from 20 (1985) to
180 94 % (2014; Table 1). Overall, however, more than 70 % of sections were surveyed in each year
181 (Table 1) and the number of missing section surveys tended to decrease with increasing distance
182 from the tidal limit and increase with increasing flow (Fig. S3).

183

184 [Table 1 here]

185

186 Linear Mixed Effects models were constructed within the statistical software package R (version
187 3.2.1, R Core Team, 2015). Models tested whether redd density varied with distance from tidal limit
188 between years, and whether this distribution of redds was affected by flow during the previous year.
189 The models compared are shown in Table 2. The saturated model was:

190

$$y \sim \alpha_i + \beta_1 Dist + \beta_2 Dist^2 + \beta_3 Flow + \beta_4 Dist \times Flow + \beta_5 Dist^2 \times Flow + \epsilon$$

191

192 where y is redd density during January/February, $Dist$ is distance from tidal limit (in km), $Flow$ is
193 either mean flows or high flows, α_i is a random intercept for year i that is Gaussian distributed with
194 mean 0 and variation σ_{year}^2 to account for variation in redd density between years that could not be
195 explained by our explanatory variables, $\beta_j \in [\beta_1, \dots, \beta_j]$ is a coefficient relating the explanatory
196 variable j to y , and ϵ is a Gaussian error term with mean 0 and variation σ^2 . We assumed a Gaussian
197 error term despite the possibility that could predict negative (<0) redd densities because it is more
198 commonly used than a truncated Gaussian distribution, such as the Gamma distribution. In practice,
199 negative redd densities were predicted only in the uppermost sections of the catchment.

200 Explanatory variables were standardised before entering the model. $Dist$ and $Flow$ were divided by
201 their standard deviation calculated for all years combined; $Dist^2$ was then calculated as the $Dist$
202 raised to the power of 2. We standardised the explanatory variables to ensure the stability of
203 coefficient estimates and thus the comparison of their effects on the response variable.

204 The models were fit by Maximum Likelihood using R package `lme4`, the estimated coefficient p
205 values were calculated using Satterthwaite's approximations implemented in the R package
206 `lmerTest`, and the conditional and marginal R squared values were calculated using the method of
207 Nakagawa & Schielzeth (2013) implemented in R package `piecewiseSEM`. There is no "best
208 practice" method to calculate standard errors for Linear Mixed Effects models and so we present
209 model fits without standard error bands.

210 Results

211 We fitted and compared five models explaining inter-annual differences in salmon redd distribution
212 along the river Frome that used linear and quadratic combinations of distance from tidal limit and
213 mean flows during the “upstream phase” of the salmon migration period. Both linear and quadratic
214 terms were included in analysis due to the curvilinear nature of redd density across the river Frome
215 (Fig. 1). Final models were chosen based on factors including Akaike Information Criteria (AIC) and
216 conditional and marginal R-squared values. Comparing the models by AIC suggested that the
217 saturated model (QuadDFint in Table 2) was the “top-ranked” model. This model included linear
218 terms for distance and mean flows and their interaction and a quadratic term for distance and its
219 interaction with flow (Table 2).

220

221 **[Table 2 here]**

222

223 The difference in AIC (dAIC) between the “top-ranked” model and the second ranked model (the
224 same model omitting the interaction between distance and flow; QuadDF) was 2.74, which exceeds
225 the oft-cited threshold value of 2 AIC points (e.g., Burnham & Anderson, 2003) indicating a
226 substantially better model, but not the more conservative threshold of 6 points (e.g., Richards,
227 2005). Moreover, the interaction between distance and mean flow in model QuadDFint was judged
228 to be statistically significant ($t_{280.19} = 1.98, p < 0.05$; Fig. 3a; Table S1) by Satterthwaite’s
229 approximation. This suggests that mean flow may affect the distribution of redds within the river
230 Frome. As this interaction was statistically significant, we retained the lower order mean flows term
231 even though it was not statistically significant (Fig. 3a). Residuals from the “top-ranked” model were
232 approximately normally distributed, suggesting that our assumption of Gaussian errors was
233 appropriate (Fig. S4).

234 Random year effect estimates generally overlapped zero (Fig. 3b). Estimates for two years, however,
235 were substantially different to zero, i.e., their standard errors did not overlap zero, suggesting that
236 the random effect was non-negligible. Moreover, the assumption that the random effect was
237 Gaussian was supported; the estimated year effects conformed to a theoretical Gaussian distribution
238 (Fig. S5).

239

240 **[Figure 3 here]**

241

242 Marginal effect plots of each term suggested a concave quadratic relationship between redd density
243 and distance (a positive effect of distance together with a negative effect of distance squared; Fig.
244 4), which was modified by a weakly convex quadratic relationship between redd density and the
245 interaction between distance and flow (a negative effect of distance by flow together with a positive
246 effect of distance squared by flow; Fig. 4). Combined, these relationships suggest that redd density
247 was highest at intermediate distances from the tidal limit under low flow and more evenly
248 distributed throughout the river during high flow. Plotting model fits to the observed redd densities
249 ordered by increasing annual critical flow appear to support this pattern (Fig. 5).

250

251 **[Figure 4 here]**252 **[Figure 5 here]**

253

254 We repeated this analysis using high flows as the measure of flow. For these models, the saturated
255 model (QuadDFint in Table 2) was again the “top-ranked” model, although the dAIC between this
256 and the second ranked model was slightly lower than for the models using mean flows as the

257 measure of flow (Supporting Information; Table S2). Unlike the “top-ranked” model for mean flow,
258 no significant relationship between high flows and redd density were observed; none of the terms
259 including flow were statistically significant by Satterthwaite’s approximation (Table S2).

260 We repeated the analyses using mean flows and high flows as the measures of flow but omitted
261 years with low survey coverage, namely years 1983, 2007, 2008 & 2012. For both measures of flow,
262 the “top-ranked” model was the model including the quadratic effect of distance and the effect of
263 flow but not including their interaction, i.e., model QuadDF (Tables S3 & S4). Again, however, the
264 dAIC between the “top-ranked” and second ranked models were lower than for the models including
265 all data, as were the marginal and conditional R^2 values, despite the additional variance present in
266 the full datasets.

267 Discussion

268 Our analysis suggests that river flow limits the upstream distance that salmon can migrate to spawn
269 and that low flows cause aggregated spawning in the middle sections of the river, with potentially
270 detrimental consequences for emerging fry subject to strong density-dependent competition (Beall
271 et al., 1994, Armstrong et al., 2003). To our knowledge, we are the first to demonstrate an inter-
272 annual effect of flow on redd distribution in a UK chalk stream. This relationship was detected
273 despite the relatively stable flow conditions in our groundwater-fed study river. The effect of flow on
274 salmon redd distribution within freshet rivers may be far greater than those experienced in chalk
275 streams, such as the river Frome.

276 Compared to years of high flow, we found that the difference in redd density between the middle
277 and upper sections of the river was accentuated in years when river flow was low during the search
278 phase of spawning migration, taken to be 1st October to 31st December on the river Frome. This was
279 characterised by a pronounced humped (or quadratic) relationship between redd density and
280 distance in the middle river sections, which might represent spawning aggregations. Conversely, we
281 found that relative to years of low flow, the difference in redd density between the middle and

282 upper sections of the river was smaller. Redds were more evenly distributed throughout the river
283 and the quadratic relationship between redd density and distance was less pronounced in years of
284 higher flow. Furthermore, when we repeated our analysis omitting years of limited survey coverage
285 due to high flows, the “top-ranked” models omitted the distance-flow interaction, providing
286 circumstantial evidence that the distance-flow interaction captured the difference in patterns of
287 redd density between high and low river flow years.

288 The increased redd density observed within the middle reaches under low flow may have occurred
289 due to two factors. Under extreme low flow, potential barriers to migration may have become
290 difficult to pass, leading to aggregations directly downstream. Salmon gathering below various weirs
291 along the river Frome has been documented since 1913 (Solomon, 2000). Aggregation below power
292 stations, weirs and other barriers have also been documented across Europe, and the ability to pass
293 these barriers within the river can be exacerbated under low flow (Thorstad et al., 2003, Ugedal et
294 al., 2008, Klett et al., 2013). Secondly, tributaries used for spawning and upstream passage might
295 become unfavourable under low flow, resulting in an increase in individuals within the middle
296 reaches of the main river. Studies on the Tay and Dee in Scotland suggest that higher water
297 discharge towards the period of final spawning migration allows salmon to access spawning
298 tributaries that are otherwise inaccessible because of insufficient water (Webb 1989, Webb &
299 Hawkins 1989). Assuming that these salmon construct redds in these otherwise inaccessible
300 tributaries, then these results would support our finding of relatively higher redd density in higher
301 river sections in high flow years. In contrast, salmon migration through the main river stem of the
302 Tay and Dee was scarcely affected by water discharge, suggesting that these deeper waters were
303 always accessible to spawning salmon (Hawkins 1989). Again assuming that these fish construct
304 redds, then these results would support our finding that redds tend to be aggregated in the middle
305 river sections, as characterised by the quadratic relationship between redd density and distance.

306 Climate change predictions forecast changing rainfall patterns across the UK, characterised by an
307 increasing mean temperature and more variable precipitation. For rivers, these predictions will likely
308 manifest in extreme flow events, including increasing incidences of droughts and floods. There are
309 many studies hypothesising how these predictions might impact salmonids in the UK and elsewhere
310 (see reviews Jonsson and Jonsson 2009; Isaak, et al. 2012). From the perspective of our findings, the
311 forecast extreme flow events and their effect on salmon spawning behaviour could have significant
312 negative impacts on spawning success. One feasible negative outcome could be large spawning
313 aggregations due to sustained low flows during the search phase of migration would result in a
314 highly competitive environment for emerging fry (Jonsson and Jonsson, 1998). Another potential
315 negative outcome might be that high flows during the search phase of migration facilitate salmon
316 entry into upstream spawning habitat that subsequently dries out under drought conditions,
317 resulting in unsuccessful redds (Becker et al., 1983, Reiser and White, 1983).

318 Any effect of flow on redd distribution might cause a shift in the behaviour and ultimately survival of
319 juvenile parr. Previous research on parr survival and movement in the river Frome revealed that
320 some parr move past the smolt monitoring station in Autumn (October – January) prior to the main
321 Spring smolt migration between March and May (Ibbotson et al., 2013). While parr that remain in
322 their natal river site have a higher probability to smoltify with increasing distance upstream, it is
323 currently unknown whether autumn migration make a significant contribution to spawning stock
324 and success (Riley et al., 2009). Under extreme low flow a higher proportion of redds might occur
325 within the middle reaches, which might lead to a higher proportion of autumn migrating parr.
326 Moreover, competition for limited resources in the middle to lower reaches of the river might
327 increase during low flow years, with detrimental consequences for individual parr survival.

328 A significant challenge in this study was to decide how best to represent flow conditions, so that
329 they accurately corresponded to the search time frame of southern UK salmon. To avoid subjectivity,
330 we followed expert opinion to define the period of final migration and summarised the flow data for

331 that period in two ways: (1) mean flows, and (2) high flows, supposed to represent mean and
332 extreme flow events, respectively. We repeated our analyses with each measure of flow separately.
333 We found that the models including the distance-flow interaction were “top-ranked” for both
334 measures of flow. The most parsimonious explanation for this finding is that the two variables were
335 highly related, i.e., were measuring the same effect. Indeed, the Pearson’s correlation coefficient
336 between the variables was > 0.9 . Nevertheless, it is reassuring that the “top-ranked” models with
337 two highly – but not perfectly – related variables included the distance-flow interaction, particularly
338 since these “top-ranked” models were not different from the second-ranked models at the more
339 conservative dAIC level of 6 (Richards 2005).

340 Although we have taken care to ensure our findings are robust, we acknowledge some study
341 limitations, several of which relate to pitfalls in redd surveying methods. Regarding the statistical
342 analysis, we used AIC for model selection, which can favour more complex models compared to, for
343 example, Bayesian Information Criteria (BIC; Burnham & Anderson, 2003). We argue that more
344 complex models will better capture the reality of redd distribution patterns. Thus reality should be
345 favoured over parsimony and AIC over BIC. Moreover, we argue that AIC should be preferred when
346 the possibility of making false negative result is considered more harmful than the possibility of
347 making a false positive result, which we consider the case with this small but valuable dataset. We
348 had a limited number of predictors to explain the inter-annual patterns of salmon redd density,
349 primarily because data on other factors are limited or have not been collected. For example,
350 locations of groundwater upwellings are believed to influence redd site selection (Saltveit, 2013) and
351 sedimentation may also affect redd success, but we do not have these data for the river Frome.

352 Aside from the statistical limitations of our findings, there are some well-known issues with redd
353 surveys that deserve mention. First, there is growing evidence that multiple males and females are
354 involved in spawning on any individual redd and that individual fish can construct multiple redds,
355 frequently $>1\text{km}$ apart (Taggart, et al. 2001). Second, redd superimposition (the construction of one

356 redd over another) was not uncommon in the Girnock Burn, a tributary of the river Dee in Scotland
357 (Taggart, et al. 2001). Finally, there are several issues that question the accuracy of redd surveys.
358 They are measured without an assessment of their associated uncertainty (e.g., Dunham et al.,
359 2001). A number of factors may affect a redd census such as the redd age, size, and colour, the
360 vegetation cover in the survey section and observer experience. Most importantly, the river depth,
361 flow and turbidity are also thought to affect the survey accuracy. To investigate the possible effect of
362 flow conditions on redd survey accuracy, we repeated our analyses omitting low survey coverage
363 years, which tended to occur in high flow years. We found that the top-ranked models for these
364 reduced data omitted the distance-flow interaction. While this could be interpreted as evidence for
365 the importance of the interaction term to describe the difference in redd distributions in low and
366 high flow years, it doesn't clarify the potential bias from poor survey accuracy because low survey
367 coverage is confounded with high flows. To better understand this potential bias requires further
368 work. In the meantime, we hold some confidence in the accuracy of the redd survey data because it
369 is often significantly correlated with spawner abundances (Beland, 1996; Dunham et al., 2001;
370 Gallagher and Gallagher, 2005).

371 While our study focused on a single UK chalk stream, and a single species, Atlantic salmon, we feel
372 that our results could generalise to other systems and fish species. There are several fish species that
373 undertake spawning migrations, including diadromous species, such as salmon and sea trout (*Salmo*
374 *trutta*) that migrate between fresh- and sea- waters, and potamodromous species, such as European
375 grayling (*Thymallus thymallus*) that migrate within the freshwater only. We see no reason why
376 migrations of these species could not also be influenced by flow conditions. Indeed, for some
377 species, such as the European grayling, the effects of flow on recruitment might be more severe
378 than for salmon because they construct comparatively shallow redds that will be less resilient to
379 extreme events which may further increase their highly variable recruitment (Crisp, 1996, Ibbotson,
380 et al. 2001). We believe our results can also be generalised to other systems, i.e., rivers. Chalk
381 streams are fed by groundwater upwellings and have stable flow regimes compared to rain-fed

382 rivers. That we were able to detect an effect of flow on redd distributions in a stable southern UK
383 chalk stream suggests that the effects of flow on the search phase of salmonid migration might be
384 more pronounced on rain-fed rivers.

385 Conclusion

386 In conclusion, this study has – for the first time - revealed the multi-annual effect of flow conditions
387 on chalk stream salmon spawning behaviour. Our results suggest that under low flow, densities
388 become aggregated within the middle reaches which could have a negative impact on juvenile
389 survival and overall recruitment. As human populations continue to grow, balancing the abstraction
390 needs of the UK with the flow requirements of salmon, all under increased climate change, is an
391 issue which will need to be addressed if we are to preserve this ecologically and economically
392 important species. Managing habitat and potentially removing weirs still present on the Frome may
393 also improve the movement of spawning salmon upstream, leading to fewer spawning aggregations
394 under extreme low flow. Our findings could also be used to revise current abstraction practices
395 during this key migration period, to improve the spawning success of salmon and other salmonids
396 under adverse flow conditions and potentially aid in future conservation of Atlantic salmon within
397 the river Frome and other chalk stream rivers.

398 Acknowledgements

399 We thank Nicole Caetano and Andy Martin of the UK Environment Agency, and Dylan Roberts,
400 William Beaumont and Luke Scott of the Game & Wildlife Conservation Trust. This work was part
401 funded through an EU Knowledge Economy Skills Scholarships grant awarded to Elinor Parry.

402 References

403 Armstrong, J. D., & Nislow, K. H. 2006. Critical habitat during the transition from maternal
404 provisioning in freshwater fish, with emphasis on Atlantic salmon (*Salmo salar*) and brown trout
405 (*Salmo trutta*). *Journal of Zoology*, 269:403-413.

- 406 Armstrong, J. D., Kemp, P. S., Kennedy, G. J. A., Ladle, M., & Milner, N. J. 2003. Habitat requirements
407 of Atlantic salmon and brown trout in rivers and streams. *Fisheries Research*, 62(2): 143-170.
- 408 Bardonnnet, A., & Baglinière, J. L. 2000. Freshwater habitat of Atlantic salmon (*Salmo salar*). *Canadian*
409 *Journal of Fisheries and Aquatic Sciences*, 57(2): 497-506.
- 410 Beall, E., Dumas, J., Claireaux, D., Barriere, L. & Marty, C., 1994. Dispersal patterns and survival of
411 Atlantic salmon (*Salmo salar* L.) juveniles in a nursery stream. *ICES Journal of Marine Science*,
412 51(1):1-9.
- 413 Becker, C. D., & Neitzel, D. A. 1985. Assessment of intergravel conditions influencing egg and alevin
414 survival during salmonid redd dewatering. *Environmental Biology of Fishes*, 12(1): 33-46.
- 415 Becker, C. D., Neitzel, D. A., & Abernethy, C. S. 1983. Effects of dewatering on Chinook salmon redds:
416 tolerance of four development phases to one-time dewatering. *North American Journal of Fisheries*
417 *Management*, 3(4): 373-382.
- 418 Beland, K. F. 1996. The relation between redd counts and Atlantic salmon (*Salmo salar*) parr
419 populations in the Dennys River, Maine. *Canadian Journal of Fisheries and Aquatic Sciences*, 53(3):
420 513-519.
- 421 Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. 2012. Impacts of climate
422 change on the future of biodiversity. *Ecology letters*, 15(4): 365-377.
- 423 Brunner, P., Dennis, I., Girvan, J. 2010 River Frome Geomorphological Assessment and Rehabilitation
424 Plan Technical Report (Draft Report). Environment Agency.
- 425 Burnham, K. P., & Anderson, D. 2003. Model selection and multi-model inference. A Practical
426 Information-Theoretic Approach. Springer.
- 427 Chapman, D. W., Weitkamp, D. E., Welsh, T. L., Dell, M. B., & Schadt, T. H. 1986. Effects of river flow
428 on the distribution of chinook salmon redds. *Transactions of the American Fisheries Society*, 115(4):
429 537-547.

- 430 Crisp, D. T., & Carling, P. A. 1989. Observations on siting, dimensions and structure of salmonid
431 redds. *Journal of Fish Biology*, 34(1): 119-134.
- 432 Crisp, D. T. 1996. Environmental requirements of common riverine European salmonid fish species in
433 fresh water with particular reference to physical and chemical aspects. *Hydrobiologia*, 323(3), 201-
434 221.
- 435 Dunham, J., Rieman, B., & Davis, K. 2001. Sources and magnitude of sampling error in redd counts
436 for bull trout. *North American Journal of Fisheries Management*, 21(2): 343-352.
- 437 Fleming, I. A. 1998. Pattern and variability in the breeding system of Atlantic salmon (*Salmo salar*),
438 with comparisons to other salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*, 55(S1): 59-
439 76.
- 440 Gallagher, S. P., & Gallagher, C. M. 2005. Discrimination of Chinook salmon, coho salmon, and
441 steelhead redds and evaluation of the use of redd data for estimating escapement in several
442 unregulated streams in northern California. *North American Journal of Fisheries Management*, 25(1):
443 284-300.
- 444 Geist, D. R., & Dauble, D. D. 1998. Redd site selection and spawning habitat use by fall chinook
445 salmon: the importance of geomorphic features in large rivers. *Environmental Management*, 22(5):
446 655-669.
- 447 Gross, M. R., Coleman, R. M., & McDowall, R. M. 1988. Aquatic productivity and the evolution of
448 diadromous fish migration. *Science*, 239(4845): 1291-1293.
- 449 Groves, P. A., & Chandler, J. A. 1999. Spawning habitat used by fall chinook salmon in the Snake
450 River. *North American Journal of Fisheries Management*, 19(4): 912-922.
- 451 Hawkins, A. D. 1989. Factors affecting the timing of entry and upstream movement of Atlantic
452 salmon in the Aberdeenshire Dee. In *Salmonid Migration and Distribution Symposium* (pp. 101-105).

- 453 Hulme, M., Jenkins, G.J., Lu, X., Turnpenny, J.R., Mitchell, T.D., Jones, R.G., Lowe, J., et al. 2002
454 Climate Change Scenarios for the United Kingdom: The UKCIP02 Scientific Report, Tyndall Centre for
455 Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich, UK.
- 456 Ibbotson, A., Cove, R., Ingraham, A., Gallagher, M., Hornby, D., Furse, M. & Williams, C. A. 2001.
457 Review of Grayling Ecology, Status and Management Practice; Recommendations for Future
458 Management in England and Wales. Environment Agency. Technical Report W245.
- 459 Ibbotson, A. T., Riley, W. D., Beaumont, W. R., C., Cook, A. C., Ives, M. J., Pinder, A. C. & Scott, L. J.
460 2013. The source of autumn and spring downstream migrating juvenile Atlantic salmon in a small
461 lowland river. *Ecology of Freshwater Fish*, 22(1): 73-81.
- 462 Isaak, D.J., Wollrab, S., Horan, D. & Chandler, G., 2012. Climate change effects on stream and river
463 temperatures across the northwest US from 1980–2009 and implications for salmonid fishes.
464 *Climatic Change*, 113(2): 499-524.
- 465 Jonsson, N., Jonsson, B., & Hansen, L. P. 1998. The relative role of density-dependent and density-
466 independent survival in the life cycle of Atlantic salmon *Salmo salar*. *Journal of Animal Ecology*,
467 67(5): 751-762.
- 468 Jonsson, B., & Jonsson, N. 2009. A review of the likely effects of climate change on anadromous
469 Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water
470 temperature and flow. *Journal of Fish Biology*, 75(10): 2381-2447.
- 471 Klett, K. J., Torgersen, C. E., Henning, J. A., & Murray, C. J. 2013. Spatial consistency of Chinook
472 Salmon redd distribution within and among years in the Cowlitz River, Washington. *North American*
473 *Journal of Fisheries Management*, 33(3), 508-518.
- 474 Kondolf, G. M., & Wolman, M. G. 1993. The sizes of salmonid spawning gravels. *Water Resources*
475 *Research*, 29(7): 2275-2285.

- 476 Lapointe, M. F., Bergeron, N. E., Bérubé, F., Pouliot, M. A., & Johnston, P. 2004. Interactive effects of
477 substrate sand and silt contents, redd-scale hydraulic gradients, and interstitial velocities on egg-to-
478 emergence survival of Atlantic salmon (*Salmo salar*). Canadian Journal of Fisheries and Aquatic
479 Sciences, 61(12): 2271-2277.
- 480 Lauridsen, R., Beaumont, W. R. C., Gregory, S., Scott, L., & Roberts, D. 2015 Fisheries Research
481 Report. Game and Wildlife Conservation Trust, 2016.
- 482 Levasseur, M., Bergeron, N. E., Lapointe, M. F., & Bérubé, F. 2006. Effects of silt and very fine sand
483 dynamics in Atlantic salmon (*Salmo salar*) redds on embryo hatching success. Canadian Journal of
484 Fisheries and Aquatic Sciences, 63(7): 1450-1459.
- 485 Mitchell, S. C., & Cunjak, R. A. 2007. Relationship of upstream migrating adult Atlantic salmon (*Salmo*
486 *salar*) and stream discharge within Catamaran Brook, New Brunswick. Canadian Journal of Fisheries
487 and Aquatic Sciences, 64:563-573.
- 488 Moir, H. J., Soulsby, C., & Youngson, A. 1998. Hydraulic and sedimentary characteristics of habitat
489 utilized by Atlantic salmon for spawning in the Girnock Burn, Scotland. Fisheries Management and
490 Ecology, 5(3): 241-254.
- 491 Moir, H. J., Soulsby, C., & Youngson, A. F. 2002. Hydraulic and sedimentary controls on the
492 availability and use of Atlantic salmon (*Salmo salar*) spawning habitat in the River Dee system, north-
493 east Scotland. Geomorphology, 45(3): 291-308.
- 494 Nakagawa, S., & Schielzeth, H. 2013. A general and simple method for obtaining R² from generalized
495 linear mixed-effects models. Methods in Ecology and Evolution, 4(2): 133-142.
- 496 National Rivers Authority South Western Region. 1995. The Frome and Piddle Catchment
497 Management Plan Consultation Report. The National Rivers Authority, UK.

- 498 Reiser, D. W., & White, R. G. 1983. Effects of complete redd dewatering on salmonid egg-hatching
499 success and development of juveniles. Transactions of the American Fisheries Society, 112(4): 532-
500 540.
- 501 Riedl, C., & Peter, A. 2013. Timing of brown trout spawning in Alpine rivers with special
502 consideration of egg burial depth. Ecology of Freshwater Fish, 22(3): 384-397.
- 503 Riley, W. D., Ibbotson, A. T. & Beaumont, W. R. C. 2009. Adult returns from Atlantic salmon, *Salmo*
504 *salar*, parr autumn migrants. Fisheries Management and Ecology, 16(1): 75-76.
- 505 Richards, S. A. 2005. Testing ecological theory using the information-theoretic approach: examples
506 and cautionary results. Ecology, 86(10): 2805-2814.
- 507 Sear, D. A., Armitage, P. D., & Dawson, F. H. 1999. Groundwater dominated rivers. Hydrological
508 Processes, 13(3): 255-276.
- 509 Solomon, D. J., Sambrook, H. T., & Broad, K. J. 1999. Salmon migration and river flow: results of
510 tracking radio tagged salmon in six rivers in south west England. Environment Agency.
- 511 Solomon, D. J. 2000. Migration Conditions on the River Frome. Environment Agency - South West
512 Region.
- 513 Solomon, D. J., & Sambrook, H. T. 2004. Effects of hot dry summers on the loss of Atlantic salmon,
514 *Salmo salar*, from estuaries in South West England. Fisheries Management and Ecology, 11(5): 353-
515 363.
- 516 Taggart, J. B., McLaren, I. S., Hay, D. W., Webb, J. H., & Youngson, A. F. 2001. Spawning success in
517 Atlantic salmon (*Salmo salar* L.): a long-term DNA profiling-based study conducted in a natural
518 stream. Molecular Ecology, 10(4): 1047-1060.

- 519 Thorstad, E. B., F. Økland, F. Kroglund & N. Jepsen, 2003. Upstream migration of Atlantic salmon at a
520 power station on the River Nidelva, Southern Norway. *Fisheries Management and Ecology* 10: 139–
521 146.
- 522 Thorstad, E. B., Økland, F., Aarestrup, K., & Heggberget, T. G. 2008. Factors affecting the within-river
523 spawning migration of Atlantic salmon, with emphasis on human impacts. *Reviews in Fish Biology*
524 and Fisheries, 18(4): 345-371.
- 525 Thuiller, W. 2007. Biodiversity: climate change and the ecologist. *Nature*, 448(7153): 550-552.
- 526 Ugedal, O., Næsje, T.F., Thorstad, E.B., Forseth, T., Saksgård, L.M. and Heggberget, T.G., 2008.
527 Twenty years of hydropower regulation in the River Alta: long-term changes in abundance of
528 juvenile and adult Atlantic salmon. *Hydrobiologia*, 609:9-23.
- 529 Warren, M., Dunbar, M. J., & Smith, C. 2015. River flow as a determinant of salmonid distribution
530 and abundance: a review. *Environmental Biology of Fishes*, 98(6): 1695-1717.
- 531 Webb, J. 1989. Movements of adult Atlantic salmon in the River Tay. *Scottish Fisheries Research*
532 Report (Vol. 44). Department of Agriculture and Fisheries for Scotland.
- 533 Webb, J., & Hawkins, A. D. 1989. Movements and spawning behaviour of adult salmon in the Girnock
534 burn, a tributary of the Aberdeenshire Dee, 1986. *Scottish Fisheries Research Report* (Vol. 40).
535 Department of Agriculture and Fisheries for Scotland.
- 536 Wessex Water Authority, 1987. Investigation into the Alleged Decline of Migratory Salmonids: Final
537 Report. Wessex Water Authority.

538 Tables

539 **Table 1.** Summary of redd, flow and survey site data each survey year, and an indication of whether
 540 it was considered a “high flow year”.


Survey Year	Mean daily flow m ³ /s		Percentage of Sites Surveyed	Total redds	High flow year	Included in analysis
	October - December	January - February				
1980-81	5.14	5.41	87	376	N	Y
1982-83	7.28	8.05	87	139	N	Y
1983-84	3.97	9.34	40	101	Y	N
1985-86	4.00	9.71	20	67	N	N
1988-89	3.57	3.65	76	308	N	Y
1991-92	5.78	3.72	73	167	N	Y
2001-02	4.04	6.40	93	335	N	Y
2004-05	4.33	4.55	47	189	N	Y
2005-06	5.91	4.25	80	345	N	Y
2007-08	5.97	8.90	37	141	Y	N
2008-09	5.58	8.24	40	237	Y	N
2009-10	6.88	8.03	67	182	N	Y
2010-11	4.34	6.86	67	129	N	Y
2011-12	2.94	3.41	93	492	N	Y
2012-13	10.82	12.67	33	60	Y	N
2014-15	4.60	6.68	94	256	N	Y

541 **Table 2.** A table presenting the Maximum Likelihood model comparison statistics comparing fits for
 542 models in which *Flow* is measured as *mean flows*: the mean daily flow from October to December in
 543 the year previous to the redd survey. The table is ordered by difference in Akaike Information
 544 Criteria (dAIC) from the “top-ranked” model (i.e., the model with the lowest AIC). Marginal and
 545 conditional R^2 were calculated according to the method of Nakagawa & Schielzeth (2013).

Model name	Model terms	Model fit			Comparison		R^2	
		Sigma	logLik	Deviance	AIC	dAIC	Marginal	Conditional
QuadDFint	<i>Dist, Dist², Flow,</i> <i>Dist × Flow,</i> <i>Dist² × Flow</i>	3.33	-775.0	1550.0	1566.0	0	0.26	0.34
QuadDF	<i>Dist, Dist², Flow</i>	3.37	-778.4	1556.8	1568.8	2.74	0.24	0.33
LinDF	<i>Dist, Flow</i>	3.73	-805.6	1611.2	1621.2	55.20	0.10	0.17
LinDFint	<i>Dist, Flow, Dist ×</i> <i>Flow</i>	3.73	-805.3	1610.6	1622.6	56.52	0.10	0.16
LinD	<i>Dist</i>	3.73	-807.4	1614.8	1622.8	56.80	0.07	0.17


546

547 Figures



548

549 **Figure 1.** Diagram showing how the density of redds is predicted to change with distance from the
550 tidal limit under low, medium and high flow conditions.




551

552 **Figure 2.** Map showing the redd survey sections on the river Frome, Dorset, UK. Major settlements


553 Wareham, Wool and Dorchester are shown in grey. Black dots indicate the start and end points of

554 each survey section. Black circles are locations of flow gauging stations East Stoke (ES) and Louden

555 Mill (LM).



556 (a)




557 (b)

558 **Figure 3.** Caterpillar plots showing Maximum Likelihood estimates of (a) the fixed effects and (b) the
 559 random effect for the “top-ranked” model. Points are the estimates; lines are the estimate standard
 560 errors; labels are the estimate values followed by an indication of their statistical significance,
 561 whereby: *** = $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$. Explanatory variable definitions are: dist_std =
 562 standardised distance; dist2_std = standardised distance squared; flow_std = standardised flow;

563 dist_flow_std = standardised distance and flow interaction; dist2_flow_std = standardised distance
564 squared and flow interaction. As flow and distance were standardised, no units are specified for
565 these variables.


For Peer Review Only



566

567 **Figure 4.** Line plots showing the marginal effects of each of the explanatory variables: `dist_std` =
 568 standardised distance; `dist2_std` = standardised distance squared; `flow_std` = standardised flow;
 569 `dist_flow_std` = standardised distance and flow interaction; `dist2_flow_std` = standardised distance
 570 squared and flow interaction. X axis is the explanatory variable value, e.g., standardised distance for
 571 the first panel. The shaded grey area is the standard error of the estimated effect. As flow and
 572 distance were standardised, no units are specified for these variables.

573



574

575 **Figure 5.** Scatter plots showing observed redd densities as a function of standardised distance from
 576 the tidal limit. Each panel represents a different year characterised by a measure of mean daily flow
 577 from 1st October to 31st December and panels are ordered from low (top left) to high (bottom right)
 578 flow. Lines are the “top-ranked” model fits. As flow and distance were standardised, no units are
 579 specified for these variables.

Supporting Information




Figure S1. Boxplot of mean monthly Atlantic salmon counts on the Frome, from 2009 to 2014, illustrating the high numbers of adults returning to the river during October – December each year to breed.






Figure S2. Scatter plots of redd density (#/km) as a function of distance from tidal limit (km) for the years that were included in the model fitting.



(a)



(b)

Figure S3. Scatter plots of number of years a section was not surveyed as a function of (a) distance from tidal limit and (b) mean daily flow. Blue lines are fitted linear regression predictions.




Figure S4. A histogram of the “top-ranked” model residuals suggesting that the assumption of normally distributed (Gaussian) errors was supported.




Figure S5. A quantile-quantile plot showing that the sample of years analysed as a random effect conform to a theoretical normal distribution, suggesting that the assumption of normally distributed year effects was supported.

Table S1. A table presenting the coefficient estimates of the top-ranking models (as assessed by AIC) when the flow predictor is either (i) *mean flows*: the mean daily flow from 1st October to 31st December or (ii) *high flows*: the number days from 1st October to 31st December that the daily flow exceeded the Q75 flow in the year previous to the redd survey.

Model name	Coefficient	Estimate (SE)	df	t	p
Mean flows	<i>Intercept</i>	1.57 (3.17)	214.26	0.50	ns
	<i>Dist</i>	20.09 (6.05)	286.50	3.32	<0.05
	<i>Dist</i> ²	-13.50 (3.00)	281.54	-4.50	<0.001
	<i>Flow</i>	-0.44 (3.54)	229.47	-0.13	ns
	<i>Dist</i> × <i>Flow</i>	-9.29 (6.54)	286.92	-1.42	ns
	<i>Dist</i> ² × <i>Flow</i>	6.22 (3.15)	280.19	1.98	<0.05
High flows	<i>Intercept</i>	1.68 (1.07)	177.95	1.57	ns
	<i>Dist</i>	12.99 (2.25)	282.65	5.77	<0.001
	<i>Dist</i> ²	-9.00 (1.22)	282.35	-7.41	<0.001
	<i>Flow</i>	-1.25 (1.50)	237.67	-0.83	ns
	<i>Dist</i> × <i>Flow</i>	-1.79 (2.81)	285.76	-0.64	ns
	<i>Dist</i> ² × <i>Flow</i>	1.78 (1.36)	281.01	1.30	ns

High flows results

Table S2. Tables of (a) Maximum Likelihood model comparison statistics comparing fits for models in which *Flow* is measured as *high flows*: the Q75 daily flow from October to December in the year previous to the redd survey. The table is ordered by difference in Akaike Information Criteria (dAIC) from the “top-ranked” model (i.e., the model with the lowest AIC). Marginal and conditional R^2 were calculated according to the method of Nakagawa & Schielzeth (2013). (b) Coefficient estimates of the top-ranking model fitted to the same data, as assessed by AIC.

(a)

Model name	Model terms	Model fit			Comparison		R^2	
		Sigma	logLik	Deviance	AIC	dAIC	Marginal	Conditional
QuadDFint	<i>Dist, Dist², Flow, Dist × Flow, Dist² × Flow</i>	3.33	-775.4	1550.8	1566.8	0.0	0.25	0.35
QuadDF	<i>Dist, Dist², Flow</i>	3.37	-778.7	1557.3	1569.3	2.5	0.23	0.33
LinDF	<i>Dist, Flow</i>	3.73	-806.1	1612.2	1622.2	55.4	0.09	0.17
LinD	<i>Dist, Flow, Dist × Flow</i>	3.73	-807.4	1614.8	1622.8	56.0	0.07	0.17
LinDFint	<i>Dist</i>	3.72	-805.8	1611.5	1623.5	56.7	0.10	0.17

(b)

Model	Coefficient	Estimate	SE	df	t	p	p level
QuadDFint	<i>Intercept</i>	1.68	1.07	177.95	1.57	0.12	ns
	<i>dist_std</i>	12.99	2.25	282.65	5.77	0.00	<0.001
	<i>dist2_std</i>	-9.00	1.22	282.35	-7.41	0.00	<0.001
	<i>flow_std</i>	-1.25	1.50	237.67	-0.83	0.41	ns
	<i>dist_flow_std</i>	-1.79	2.81	285.76	-0.64	0.52	ns
	<i>dist2_flow_std</i>	1.78	1.36	281.01	1.30	0.19	ns

Removing “odd” years

Mean flows results

Table S3. Tables of (a) Maximum Likelihood model comparison statistics comparing fits for models in which *Flow* is measured as *high flows*: the Q75 daily flow from October to December in the year previous to the redd survey and years with low survey coverage (1983, 2007, 2008 & 2012) have been removed. The table is ordered by difference in Akaike Information Criteria (dAIC) from the “top-ranked” model (i.e., the model with the lowest AIC). Marginal and conditional R^2 were calculated according to the method of Nakagawa & Schielzeth (2013). (b) Coefficient estimates of the top-ranking model fitted to the same data, as assessed by AIC.

(a)

Model name	Model terms	Model fit			Comparison		R^2	
		Sigma	logLik	Deviance	AIC	dAIC	Marginal	Conditional
QuadDF	<i>Dist, Dist², Flow</i>	3.49	-655.79	1311.6	1323.6	0.00	0.27	0.31
QuadDFint	<i>Dist, Dist², Flow, Dist × Flow, Dist² × Flow</i>	3.46	-654.09	1308.2	1324.2	0.62	0.28	0.32
LinDF	<i>Dist, Flow</i>	3.88	-679.59	1359.2	1369.2	45.60	0.12	0.15
LinDFint	<i>Dist, Flow, Dist × Flow</i>	3.87	-679.15	1358.3	1370.3	46.73	0.12	0.15
LinD	<i>Dist</i>	3.87	-682.78	1365.6	1373.6	49.99	0.07	0.14

(b)

Model	Coefficient	Estimate	SE	df	t	p	p level
QuadDF	<i>Intercept</i>	5.36	1.55	18.44	3.47	0.00	<0.05
QuadDF	<i>dist_std</i>	11.46	1.99	236.58	5.75	0.00	<0.001
QuadDF	<i>dist2_std</i>	-7.92	1.09	236.61	-7.29	0.00	<0.001
QuadDF	<i>flow_std</i>	-4.46	1.49	10.66	-2.99	0.01	<0.05

High flows results

Table S4. Tables of (a) Maximum Likelihood model comparison statistics comparing fits for models in which *Flow* is measured as *high flows*: the Q75 daily flow from October to December in the year previous to the redd survey and years with low survey coverage (1983, 2007, 2008 & 2012) have been removed. The table is ordered by difference in Akaike Information Criteria (dAIC) from the “top-ranked” model (i.e., the model with the lowest AIC). Marginal and conditional R^2 were calculated according to the method of Nakagawa & Schielzeth (2013). (b) Coefficient estimates of the top-ranking model fitted to the same data, as assessed by AIC.

(a)

Model name	Model terms	Model fit			Comparison		R^2	
		Sigma	logLik	Deviance	AIC	dAIC	Marginal	Conditional
QuadDF	<i>Dist, Dist², Flow</i>	3.48	-657.73	1315.47	1327.47	0.00	0.24	0.32
QuadDFint	<i>Dist, Dist², Flow, Dist × Flow, Dist² × Flow</i>	3.47	-656.47	1312.93	1328.93	1.47	0.25	0.32
LinDF	<i>Dist, Flow</i>	3.87	-681.58	1363.16	1373.16	45.69	0.09	0.15
LinD	<i>Dist</i>	3.87	-682.78	1365.56	1373.56	46.09	0.07	0.14
LinDFint	<i>Dist, Flow, Dist × Flow</i>	3.87	-681.13	1362.26	1374.26	46.80	0.09	0.15

(b)

Model	Coefficient	Estimate	SE	df	t	p	p level
QuadDF	<i>Intercept</i>	1.95	0.96	101.44	2.02	0.05	<0.05
QuadDF	<i>dist_std</i>	11.52	1.99	236.28	5.78	0.00	<0.001
QuadDF	<i>dist2_std</i>	-7.93	1.09	236.41	-7.31	0.00	<0.001
QuadDF	<i>flow_std</i>	-1.17	0.72	11.95	-1.61	0.13	ns

For Peer Review Only

Table S5. Survey section details including survey section length (calculated using GIS) and distance from tidal limit. The location of the start and end points of each survey section are illustrated in figure 2.

Survey Section Number and Name	Section Length (in kilometres)	Start point distance from tidal limit (in kilometres)
1 Worgret - Holme Bridge	7.78	0
2 Holme Bridge - East Stoke Weir	3.50	7.80
15 East Stoke Mill Stream	1.0	11.80
3 East Stoke Weir - Bindon Hatches	2.50	14.70
16 Bindon Mill Stream	0.25	18.50
24 Wool Stream	1.75	18.70
4 Bindon Hatches - East Burton Hatches	4.50	19.20
17 Waterbarn Stream	1.75	21.80
19 Moreton North Stream	3.25	23.90
18 Trout Stream	0.50	26.30
5 E Burton Hatches - Moreton House Weir	1.75	28.0
25 Tadnoll Brook	8.0	28.80
6 Moreton House Weir - Hurst Weir	1.75	31.30
7 Hurst Weir - Woodsford Weir	4.50	34.90
20 North Stream	5.0	39.50
8 Woodsford Weir - Nine Hatches	1.25	43.0
9 Nine Hatches - Stafford House Weir	3.0	45.70
20a North Stream south arm	1.50	48.60
21 Greys Bridge Carrier	2.50	50.0
10 Stafford House Weir - Louds Mill	2.75	51.50
26 South Winterbourne	8.50	51.50
22 South Winterbourne Carrier	2.25	55.0
11 Louds Mill - Whitfield Hatches	3.75	56.60
23a Dorchester Mill Stream	1.41	57.70
23 North Dorchester Carrier	6.0	60.40
27 River Cerne	14.50	63.10
12 Whitfield Hatches - Frampton Weir	4.75	63.20
28 Sydling Brook	72.10	72.10
13 Frampton Weir - Hooke confluence	6.25	74.70
14 Hooke confluence - Cattistock	1.75	83.70
29 River Hooke	3.62	83.70