

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/98437/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Riglin, Lucy, Collishaw, Stephan, Richards, Alexander, Thapar, Ajay, Maughan, Barbara, O'Donovan, Michael Conlon and Thapar, Anita 2017. Schizophrenia polygenic risk score and psychotic risk detection - authors' reply. *The Lancet Psychiatry* 4 (3) , pp. 188-189. 10.1016/S2215-0366(17)30052-4 file

Publishers page: [http://dx.doi.org/10.1016/S2215-0366\(17\)30052-4](http://dx.doi.org/10.1016/S2215-0366(17)30052-4)  
< [http://dx.doi.org/10.1016/S2215-0366\(17\)30052-4](http://dx.doi.org/10.1016/S2215-0366(17)30052-4) >

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## Schizophrenia polygenic risk score and psychotic risk detection – authors' reply

Lucy Riglin, PhD<sup>1</sup>

Stephan Collishaw, PhD<sup>1</sup>

Alexander Richards, PhD<sup>1</sup>

Ajay K Thapar, MRCPGP<sup>1</sup>

Barbara Maughan, PhD<sup>2</sup>

Michael C O'Donovan, FRCPsych<sup>1</sup>

Anita Thapar, FRCPsych<sup>1\*</sup>

<sup>1</sup>Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, UK; <sup>2</sup>MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK. \*Corresponding author. Division of Psychological Medicine and Clinical Neurosciences, Cardiff University School of Medicine, Hadyr Ellis Building, Maindy Road, Cathays, Cardiff CF24 4HQ. Tel: +442920 6688325. Email: [thapar@cardiff.ac.uk](mailto:thapar@cardiff.ac.uk).

We thank Michele Poletti and colleagues for their interest in our recent population study in which we investigated association between schizophrenia risk alleles and neurodevelopmental outcomes in childhood.<sup>1</sup> We agree that child and adolescent mental health services (CAMHS) are a good place to detect individuals at elevated risk for schizophrenia. We also agree that a developmentally-informed framework for risk assessment is important. Given that disorders such as schizophrenia can manifest as neurodevelopmental impairments earlier in life<sup>2,3</sup> (which may index genetic liability<sup>1</sup>), conceptualizing early risk as only the presentation of psychotic-like features in adolescence and adult life is likely to be limiting. However, the majority of impairments found to precede schizophrenia are non-specific; cognitive, social, behavioural and emotional impairments have also been found to precede other psychiatric disorders, such as depression.<sup>2,3</sup> Thus distinguishing individuals who are specifically at risk for psychosis is complex and there will be challenges in stratifying risk appropriately. Nevertheless, for children already at elevated risk of developing schizophrenia (e.g. who have a parent with the disorder), increased attention to early neurodevelopmental impairments may aid identification of those at ultra-high risk. Findings from our study and the work of others<sup>4</sup> suggest that before the typical age of illness onset, schizophrenia (genetic) liability may manifest as symptoms that do not resemble psychosis. Identifying those at risk in CAMHS is only of benefit if services are able to provide interventions that reduce risk but that is another challenge. Moreover accessing CAMHS services remains a major limitation in many countries. As we highlight in our paper<sup>1</sup> intervention studies that target early impairments would be useful to differentiate between impairments that are causally associated with later schizophrenia and those that reflect pleiotropy. However much more work is needed to identify interventions that are effective in reducing risk of future psychosis. Nevertheless, we agree with Poletti and colleagues - a first step towards this is to adopt a developmental conceptualization of schizophrenia.

We declare no competing interests.

## References

1. Riglin L, Collishaw S, Richards A, et al. Schizophrenia risk alleles and neurodevelopmental outcomes in childhood: a population-based cohort study. *Lancet Psychiatry*. 2017;4(1):57-62.
2. Rutter M, Kim-Cohen J, Maughan B. Continuities and discontinuities in psychopathology between childhood and adult life. *Journal of Child Psychology and Psychiatry*. Mar-Apr 2006;47(3-4):276-295.
3. Pine DS, Fox NA. Childhood antecedents and risk for adult mental disorders. *Annual Review of Psychology*. 2015;66:459-485.
4. Jones HJ, Stergiakouli E, Tansey KE, et al. Phenotypic manifestation of genetic risk for schizophrenia during adolescence in the general population. *JAMA Psychiatry*. Published online January 27, 2016 2016.