

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/98565/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Song, Peng, Deng, Bailin , Wang, Ziqi, Dong, Zhichao, Li, Wei, Fu, Chi-Wing and Liu, Ligang 2016. CofiFab: coarse-to-fine fabrication of large 3D objects. ACM Transactions on Graphics 35 (4) , 45. 10.1145/2897824.2925876

Publishers page: <http://dx.doi.org/10.1145/2897824.2925876>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Supplementary Material for CofiFab: Coarse-to-Fine Fabrication of Large 3D Objects

1 Volumes of convex polyhedrons

To compute the volume $V(\mathbf{P})$ for a convex polyhedron P with vertices $\mathbf{P} = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$, we first introduce a new vertex

$$\mathbf{p}(f_j) = \frac{1}{\sigma(j)} \sum_{k=1}^{\sigma(j)} \mathbf{p}_{j_k}$$

for every non-triangular face f_j with vertices $\mathbf{p}_{j_1}, \mathbf{p}_{j_2}, \dots, \mathbf{p}_{j_{\sigma(j)}}$. Connecting $\mathbf{p}(f_j)$ with all vertices of f_j results in a triangulation of the polyhedron. Then the volume of the polyhedron can be computed as [Allgower and Schmidt 1986]

$$V(\mathbf{P}) = \frac{1}{6} \sum_{t_i \in \mathcal{T}} \det(\mathbf{p}^1(t_i), \mathbf{p}^2(t_i), \mathbf{p}^3(t_i)), \quad (1)$$

where \mathcal{T} is the set of faces for the triangulated polyhedron, and $\mathbf{p}^1(t_i), \mathbf{p}^2(t_i), \mathbf{p}^3(t_i)$ are the vertices of triangle t_i in positive orientation. In our optimization, the positive orientation is determined from the initial polyhedron shape, by choosing a consistent ordering of triangle vertices such that Equation (1) produces a positive value.

2 Surface sampling for convex polyhedrons

Our optimization requires sample points $\{\mathbf{q}_i\}$ on the surface of a polyhedron P , represented as $\mathbf{q}_i = \mathbf{P}\mathbf{b}_i$, where $\mathbf{b}_i \in \mathbb{R}^n$ are pre-computed convex combination coefficients with respect to the polyhedron vertex positions. To generate the samples and compute the coefficient vectors $\{\mathbf{b}_i\}$, we first triangulate the polyhedron by introducing new vertices on non-triangular faces (see Section 1). We then compute three types of sample points from the triangulated polyhedron T :

1. *Vertices of T* : such a sample point \mathbf{q}_i is either a vertex of the original polyhedron P , or an interior point on a face of P . In the former case, vector \mathbf{b}_i has exactly one non-zero element of value 1. In the latter case, there are $\sigma(j)$ non-zero elements in \mathbf{b}_i , each with value $1/\sigma(j)$, where $\sigma(j)$ is the number of vertices of the original polyhedron face that contains \mathbf{q}_i (see Equation (1)).
2. *Interior points on an edge e_i of T* : such a point can be represented as a convex combination of the two vertex sample points that belongs to e_i . In our implementation, we generate K internal sample points for each edge. Let $\mathbf{q}_{i_1}, \mathbf{q}_{i_2}$ be the coefficient vectors for the two end vertex samples for e_i , then the K interior samples on e_i are computed as:

$$\mathbf{q}_j(e_i) = \frac{j}{K+1} \mathbf{q}_{i_1} + \frac{K-j+1}{K+1} \mathbf{q}_{i_2}, \quad j = 1, \dots, K.$$

3. *Interior points on a triangle t_i of T* : such a point can be represented as a convex combination of the three vertex sample points that belongs to t_i . Let $\mathbf{q}_{i_1}, \mathbf{q}_{i_2}, \mathbf{q}_{i_3}$ be the coefficient vectors for the vertex samples, then according to the parameter K the sample points are computed as:

$$\mathbf{q}_{a,b,c}(t_i) = \frac{a}{K+1} \mathbf{q}_{i_1} + \frac{b}{K+1} \mathbf{q}_{i_2} + \frac{c}{K+1} \mathbf{q}_{i_3},$$

where $a, b, c \in \mathbb{N}$ and $a + b + c = K + 1$.

We determine the value of K from a user-specified parameter N_s for the preferred number of samples. K is chosen as the smallest number such that the total number of sample points is at least N_s .

3 Computation of centroids

To compute the centroid \mathbf{C} of the final model, we consider the final model as the combination of a hollow polyhedron made from uniform thin-sheet materials, and a 3D volume shell with uniform density. Then

$$\mathbf{C} = \frac{(\mathbf{C}_1 V_1 - \mathbf{C}_3 V_3) \rho_1 + \mathbf{C}_2 A_2 \rho_2}{(V_1 - V_3) \rho_1 + A_2 \rho_2},$$

where $\mathbf{C}_1, \mathbf{C}_3$ are the solid centroids of the target shape and the polyhedron, respectively; \mathbf{C}_2 is the surface centroid of the polyhedron; V_1, V_3 are the internal volumes of the target surface and the polyhedron, respectively; A_2 is the polyhedron surface area; ρ_1 and ρ_2 are parameters for the volume density of the 3D printed part and the area density of the laser-cut material, respectively. Here V_1, V_3 can be computed using Equation (1). Using the same notation as Equation (1), the solid centroid of a polyhedron shape can be computed as

$$\mathbf{C}(\mathbf{P}) = \frac{\sum_{t_i \in \mathcal{T}} \det(\mathbf{p}^1(t_i), \mathbf{p}^2(t_i), \mathbf{p}^3(t_i)) (\mathbf{p}^1(t_i) + \mathbf{p}^2(t_i) + \mathbf{p}^3(t_i))}{4 \cdot \sum_{t_i \in \mathcal{T}} \det(\mathbf{p}^1(t_i), \mathbf{p}^2(t_i), \mathbf{p}^3(t_i))}, \quad (2)$$

while the surface area of a polyhedron is

$$A_{\mathbf{P}} = \frac{1}{2} \sum_{t_i \in \mathcal{T}} \|[\mathbf{p}^2(t_i) - \mathbf{p}^1(t_i)] \times [\mathbf{p}^3(t_i) - \mathbf{p}^1(t_i)]\|, \quad (3)$$

and its surface centroid is

$$\mathbf{C}_A(\mathbf{P}) = \frac{\sum_{t_i \in \mathcal{T}} \|[\mathbf{p}^2(t_i) - \mathbf{p}^1(t_i)] \times [\mathbf{p}^3(t_i) - \mathbf{p}^1(t_i)]\| \sum_{k=1}^3 \mathbf{p}^k(t_i)}{\sum_{t_i \in \mathcal{T}} \|[\mathbf{p}^2(t_i) - \mathbf{p}^1(t_i)] \times [\mathbf{p}^3(t_i) - \mathbf{p}^1(t_i)]\|}, \quad (4)$$

$\mathbf{C}_1, \mathbf{C}_3$ are computed using formula (2), while A_s and \mathbf{C}_2 are computed using formulas (3) and (4), respectively.

4 Constraints for optimizing multiple polyhedrons

The two faces (f_k^i, f_l^j) chosen for the connection between two polyhedrons must satisfy the following conditions:

1. f_k^i, f_l^j are parallel, with their outward normals pointing towards each other;
2. there exists a cylinder with radius r and with its axis parallel to the normals of f_k^i, f_l^j , such that its two ends touch the two faces (f_k^i, f_l^j) and lie within the interior of each face, and the whole cylinder lie inside the target shape.

For the first condition, we require

$$\mathbf{n}_k^i + \mathbf{n}_l^j = \mathbf{0},$$

where \mathbf{n}_k^i and \mathbf{n}_l^j are the outward normal variables for the two faces. For the second condition, we introduce auxiliary variables $\mathbf{c}_k^i, \mathbf{c}_l^j \in \mathbb{R}^3$ for the centers of the circles, where the cylinder touches the two faces. \mathbf{c}_k^i and \mathbf{c}_l^j are required to lie on the two faces, respectively. The line segment between these two points must be orthogonal to the two faces, thus requiring

$$\mathbf{c}_k^i + t_k^i \mathbf{n}_k^i = \mathbf{c}_l^j,$$

with auxiliary variable $t_k^i > 0$. Moreover, each face must be kept inside a disc with radius r and center \mathbf{c}_k^i (or \mathbf{c}_l^j , respectively). Taking face f_k^i as an example, we require

$$(\mathbf{c}_k^i - \mathbf{p}_{j_1}) \cdot \frac{\mathbf{n}_k^i \times (\mathbf{p}_{j_1} - \mathbf{p}_{j_2})}{\|\mathbf{n}_k^i \times (\mathbf{p}_{j_1} - \mathbf{p}_{j_2})\|} \geq r,$$

where $\mathbf{p}_{j_1}, \mathbf{p}_{j_2}$ are two adjacent vertices in f_k^i in an appropriate order. A similar constraint is defined for face f_l^j . Finally, we compute a set of sample points $\{\mathbf{q}\}$ on the cylinder, and enforce a constraint

$$D(\mathbf{q}) \geq d_{\min},$$

where D is the signed distance function from the surface of the whole object. Each sample \mathbf{q} is computed as

$$\mathbf{q} = a\mathbf{c}_k^i + (1-a)\mathbf{n}_l^j + r(\mathbf{e}_1^{k,i} \cos b + \mathbf{e}_2^{k,i} \sin b),$$

where parameters $a \in [0, 1]$ and $b \in [0, 2\pi]$ are pre-determined, $\mathbf{e}_1^{k,i}, \mathbf{e}_2^{k,i}$ are auxiliary variables that form an orthonormal frame with \mathbf{n}_k^i , previously used for enforcing the bounding rectangle constraints.

References

ALLGOWER, E. L., AND SCHMIDT, P. H. 1986. Computing volumes of polyhedra. *Math. Comput.* 46, 173, 171–174.