


Open Access

Gamal A. El-Hiti*, Keith Smith, Amany S. Hegazy and Benson M. Kariuki

Crystal structure of 1,1-dimethyl-3-(4-methylphenyl)urea, C₁₀H₁₄N₂O



DOI 10.1515/ncls-2016-0290

Received September 30, 2016; accepted January 13, 2017; available online February 21, 2017

Abstract

C₁₀H₁₄N₂O, monoclinic, Cc (no. 9), $a = 6.8354(9)$ Å, $b = 13.8221(17)$ Å, $c = 10.2905(12)$ Å, $\beta = 90.625(11)^\circ$, $V = 972.2(2)$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0526$, $wR_{\text{ref}}(F^2) = 0.1335$, $T = 296(2)$ K.

CCDC no.: 1527333

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

The title compound was synthesised by reaction of equimolar quantities of 4-methylaniline and dimethylcarbamoyl chloride in dichloromethane at 40 °C for 1 h in the presence of excess triethylamine (1.38 mole equivalents). The crude product was recrystallized from a 1:3 mixture (by volume) of ethyl acetate and diethyl ether to give 1,1-dimethyl-3-(4-methylphenyl)urea (92%) as colourless crystals, Mp.: 141–142 °C (lit. 137–138 °C [1]; 152–153 °C [2]).

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Table 1: Data collection and handling.

| | |
|--|---|
| Crystal: | Colourless plate |
| Size: | 0.45 × 0.35 × 0.10 mm |
| Wavelength: | Mo K α radiation (0.71073 Å) |
| μ : | 0.8 cm ⁻¹ |
| Diffractometer, scan mode: | SuperNova, ω -scans |
| 2 θ_{max} , completeness: | 59.6°, >99% |
| $N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} : | 3791, 3791, 0.059 |
| Criterion for I_{obs} , $N(hkl)_{\text{gt}}$: | $I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2943 |
| $N(\text{param})_{\text{refined}}$: | 120 |
| Programs: | CrysAlis ^{PRO} [17], SHELX [18], PLATON [19] |

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

| Atom | x | y | z | $U_{\text{iso}}^{*}/U_{\text{eq}}$ |
|------|------------|-----------|-----------|------------------------------------|
| C1 | 0.0013(4) | 0.6008(2) | 0.5771(3) | 0.0391(7) |
| C2 | -0.1240(5) | 0.6047(3) | 0.4684(4) | 0.0517(10) |
| H2 | -0.0968 | 0.5687 | 0.3945 | 0.062* |
| C3 | -0.2885(5) | 0.6629(3) | 0.4726(4) | 0.0584(11) |
| H3 | -0.3714 | 0.6651 | 0.4004 | 0.070* |
| C4 | -0.3341(4) | 0.7180(3) | 0.5803(4) | 0.0489(9) |
| C5 | -0.2091(4) | 0.7138(3) | 0.6860(4) | 0.0489(8) |
| H5 | -0.2366 | 0.7502 | 0.7596 | 0.059* |
| C6 | -0.0424(4) | 0.6559(3) | 0.6848(4) | 0.0452(8) |
| H6 | 0.0402 | 0.6543 | 0.7571 | 0.054* |
| C7 | -0.5184(5) | 0.7786(3) | 0.5819(6) | 0.0691(13) |
| H7A | -0.6305 | 0.7372 | 0.5721 | 0.104* |
| H7B | -0.5158 | 0.8242 | 0.5117 | 0.104* |
| H7C | -0.5259 | 0.8126 | 0.6630 | 0.104* |
| C8 | 0.2587(4) | 0.4929(2) | 0.4877(4) | 0.0392(7) |
| C9 | 0.5152(6) | 0.3823(3) | 0.4257(5) | 0.0587(10) |
| H9A | 0.4308 | 0.3757 | 0.3511 | 0.088* |
| H9B | 0.5441 | 0.3195 | 0.4608 | 0.088* |
| H9C | 0.6347 | 0.4134 | 0.4007 | 0.088* |
| C10 | 0.5180(5) | 0.4470(4) | 0.6474(5) | 0.0591(11) |
| H10A | 0.6537 | 0.4311 | 0.6369 | 0.089* |
| H10B | 0.4598 | 0.4025 | 0.7072 | 0.089* |
| H10C | 0.5068 | 0.5116 | 0.6805 | 0.089* |
| N1 | 0.1702(4) | 0.5411(2) | 0.5869(3) | 0.0422(6) |
| H1 | 0.2219 | 0.5347 | 0.6629 | 0.051* |
| N2 | 0.4186(4) | 0.4404(2) | 0.5234(4) | 0.0490(7) |
| O1 | 0.1970(4) | 0.4948(2) | 0.3756(3) | 0.0590(8) |

Experimental details

The crystal was twinned and the structure was refined using option HKL 5 of the SHELX program [18]. All hydrogen atoms were placed in calculated positions and refined using a riding model. Methyl C-H bonds were fixed at 0.96 Å, with displacement parameters 1.5 times $U_{\text{eq}}(\text{C})$, and were allowed to spin about the C-C bond. Aromatic C-H distances were set to 0.93 Å and N-H to 0.86 Å and their U_{iso} set to 1.2 times the U_{eq} for the atoms to which they are bonded.

Discussion

Urea derivatives are important intermediates in medicinal chemistry [3–5]. Many convenient processes have been reported for the production of substituted ureas [6–11]. Aromatic ureas can be substituted efficiently *via* lithiation chemistry [12–16].

The asymmetric unit consists of one molecule of C₁₀H₁₄N₂O. The molecule is not planar as indicated by torsion angles C2—C1—N1—C8 = 13.1(5)° and N1—C8—N2—C10 = −12.2(5)°. N—H···O hydrogen bonds form chains of molecules along [001].

Acknowledgements: The project was supported by King Saud University, Deanship of Scientific Research, Research Chair.

References

1. Hutchby, M.: Novel Synthetic Chemistry of Ureas and Amides. Springer Thesis, Springer-Verlag, Berlin, Heidelberg, (2013).
2. Franz, R. A.; Applegath, F.; Morriss, F. V.; Breed, L. W.: The preparation of unsymmetrical ureas from carbon monoxide, sulfur, and amines. *J. Org. Chem.* **27** (1962) 4341–4346.
3. Kozikowski, A. P.; Zhang, J.; Nan, F.; Petukhov, P.; Grajkowska, A. E.; Wroblewski, J. T.; Yamamoto, T.; Bzdega, T.; Wroblewska, B.; Neale, J. H.: Synthesis of urea-based inhibitors as active site probes of glutamate carboxypeptidase II: efficacy as analgesic agents. *J. Med. Chem.* **47** (2004) 1729–1738.
4. dos Santos, L.; Lima, L. A.; Cechinel-Filho, V.; Correa, R.; de Campos Buzzi, F.; Nunes, R. J.: Synthesis of new 1-phenyl-3-{4-[*(2E*)-3-phenylprop-2-enoyl]phenyl}-thiourea and urea derivatives with anti-nociceptive activity. *Bioorg. Med. Chem.* **16** (2008) 8526–8534.
5. Khan, K. M.; Saeed, S.; Ali, M.; Gohar, M.; Zaid, J.; Khan, A.; Pervaen, S.; Choudhary, M. I.: Unsymmetrically disubstituted urea derivatives: a potent class of antiglycating agents. *Bioorg. Med. Chem.* **17** (2009) 2447–2451.
6. Artuso, E.; Degani, I.; Fochi, R.; Magistris, C.: Preparation of mono-, di-, and trisubstituted ureas by carbonylation of aliphatic amines with *S,S*-dimethyl dithiocarbonate. *Synthesis* **39** (2007) 3497–3506.
7. Mizuno, T.; Nakai, T.; Mihara, M.: Synthesis of unsymmetrical ureas by sulfur-assisted carbonylation with carbon monoxide and oxidation with molecular oxygen under mild conditions. *Synthesis* **41** (2009) 2492–2496.
8. Wu, C.; Cheng, H.; Liu, R.; Wang, Q.; Hao, Y.; Yu, Y.; Zhao, F.: Synthesis of urea derivatives from amines and CO₂ in the absence of catalyst and. Solvent. *Green Chem.* **12** (2010) 1811–1816.
9. Vinogradova, E. V.; Fors, B. P.; Buchwald, S. L.: Palladium-catalyzed cross-coupling of aryl chlorides and triflates with sodium cyanate: a practical synthesis of unsymmetrical ureas. *J. Am. Chem. Soc.* **134** (2012) 11132–11135.
10. Carnaroglio, D.; Martina, K.; Palmisano, G.; Penoni, A.; Domini, C.; Cravotto, G.: One-pot sequential synthesis of isocyanates and urea derivatives via a microwave-assisted Staudinger–aza-Wittig reaction. *Beilstein J. Org. Chem.* **9** (2013) 2378–2386.
11. Thalluri, K.; Manne, S. R.; Dev, D.; Mandal, B.: Ethyl 2-cyano-2-(4-nitrophenylsulfonyloxyimino)acetate-mediated Lossen rearrangement: single-pot racemization-free synthesis of hydroxamic acids and ureas from carboxylic acids. *J. Org. Chem.* **79** (2014) 3765–3775.
12. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Directed lithiation of *N'*-(2-(4-methoxyphenyl)ethyl)-*N,N*-dimethylurea and *tert*-butyl (2-(4-methoxyphenyl)ethyl)carbamate. *Synthesis* **46** (2014) 394–402.
13. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Control of site of lithiation of 3-(aminomethyl)pyridine derivatives. *Synthesis* **45** (2013) 3426–3434.
14. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Variation in the site of lithiation of 2-(2-methylphenyl)ethanamine derivatives. *J. Org. Chem.* **77** (2012) 11210–11215.
15. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Lithiation and substitution of *N'*-(ω -phenylalkyl)-*N,N*-dimethylureas. *Synthesis* **44** (2012) 20139–2022.
16. Smith, K.; El-Hiti, G. A.; Hegazy, A. S.: Lateral lithiation of *N'*-(2-methylbenzyl)-*N,N*-dimethylurea and *N*-(2-methylbenzyl)pivalamide: Synthesis of tetrahydroisoquinolines. *Synthesis* **42** (2010) 1371–1380.
17. Agilent Technologies: CrysAlis^{PRO} Software system, Agilent Technologies, Yarnton, England (2014).
18. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
19. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.